

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number  
**WO 02/24672 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 307/00**

[US/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **WUSTROW, David, Juergen** [US/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US).

(21) International Application Number: PCT/US01/25892

(22) International Filing Date: 17 August 2001 (17.08.2001)

(74) Agents: **RICHARDS, John** et al.; Ladas & Parry, 26 West 61st Street, New York, NY 10023 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/233,786 19 September 2000 (19.09.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*):  
**WARNER-LAMBERT COMPANY** [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BORA, Keenan, Martin** [US/US]; 541 Elizabeth Apt. #2, Ann Arbor, MI 48104 (US). **HU, Lain-Yen** [—/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **KESTEN, Suzanne, Ross** [US/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **LEI, Huanyshu** [CA/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **MORELAND, David, Winslow** [US/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **RAFFERTY, Michael, Francis** [US/US]; 2800 Plymouth Road, Ann Arbor, MI 48105 (US). **RYDER, Todd, Robert** [US/US]; 91 Cornhill Place, Rochester, NY 14608 (US). **SCHOLTEN, Jeffrey, David**

Published:

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: BRANCHED CHAIN AMINO ACID-DEPENDENT AMINOTRANSFERASE INHIBITORS AND THEIR USE IN THE TREATMENT OF NEURODEGENERATIVE DISEASES

(57) Abstract: The invention relates to BCAT inhibitors and use thereof for treating or preventing neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycemia and surgery, as well as treating neurodegenerative diseases including Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease and Down's syndrome, treating or preventing the adverse consequences of the overstimulation of the excitatory amino acids, treating anxiety, psychosis, convulsions, aminoglycoside antibiotics-induced hearing loss, migraine headache, chronic pain, neuropathic pain, Parkinson's disease, diabetic retinopathy, glaucoma, CMV retinitis, urinary incontinence, opioid tolerance or withdrawal, and inducing anesthesia, as well as for enhancing cognition.



WO 02/24672 A2

- 1 -

BRANCHED CHAIN AMINO ACID-DEPENDENT AMINOTRANSFERASE  
INHIBITORS AND THEIR USE IN THE TREATMENT OF  
NEURODEGENERATIVE DISEASES

FIELD OF THE INVENTION

5           This invention is related to branched chain amino acid-dependent amino  
transferase (BCAT) inhibitors. The invention is also directed to the use of BCAT  
inhibitors as neuro-protective agents for treating conditions such as stroke, cerebral  
ischemia, central nervous system trauma, hypoglycemia, anxiety, convulsions,  
aminoglycoside antibiotics-induced hearing loss, migraine headaches, chronic pain,  
10   neuropathic pain, glaucoma, CMV retinitis, diabetic retinopathy, psychosis, urinary  
incontinence, opioid tolerance or withdrawal, or neuro-degenerative disorders such as  
lathyrism, Alzheimer's disease, Parkinsonism, amyotrophic lateral sclerosis (ALS),  
and Huntington's Disease.

RELATED BACKGROUND ART

15           Excessive excitation by neurotransmitters can cause the degeneration and death  
of neurons. It is believed that this degeneration is in part mediated by the excitotoxic  
actions of the excitatory amino acids (EAA) glutamate and aspartate at the N-methyl-  
D-aspartate (NMDA) receptor. This excitotoxic action is considered responsible for  
the loss of neurons in cerebrovascular disorders such as cerebral ischemia or cerebral  
20   infarction resulting from a range of conditions, such as thromboembolic or  
hemorrhagic stroke, cerebral vasospasms, hypoglycemia, cardiac arrest, status  
epilepticus, perinatal asphyxia, anoxia such as from drowning, pulmonary surgery and  
cerebral trauma, as well as lathyrism, Alzheimer's disease, Parkinson's disease, and  
Huntington's disease.

25           Excitatory amino acid receptor antagonists that block NMDA receptors are  
recognized for usefulness in the treatment of disorders. NMDA receptors are  
intimately involved in the phenomenon of excitotoxicity, which may be a critical

- 2 -

determinant of outcome of several neurological disorders. Disorders known to be responsive to blockade of the NMDA receptor include acute cerebral ischemia (stroke or cerebral trauma, for example), muscular spasm, convulsive disorders, neuropathic pain and anxiety, and may be a significant causal factor in chronic neurodegenerative disorders such as Parkinson's disease (Klockgether T., Turski L., *Ann. Neurol.*, 1993;34:585-593), human immunodeficiency virus (HIV) related neuronal injury, amyotrophic lateral sclerosis (ALS), Alzheimer's disease (Francis P.T., Sims N.R., Procter A.W., Bowen D.M., *J. Neurochem.*, 1993;60(5):1589-1604, and Huntington's disease (see Lipton S., *TINS*, 1993;16(12):527-532; Lipton S.A., Rosenberg P.A., *New Eng. J. Med.*, 1994;330(9):613-622; and Bigge C.F., *Biochem. Pharmacol.*, 1993;45:1547-1561, and referenced cited therein.) NMDA receptor antagonists may also be used to prevent tolerance to opiate analgesia or to help control withdrawal symptoms from addictive drugs (Eur. Pat. Appl. 488,959A).


U.S. Pat. No. 5,352,683 discloses the treatment of chronic pain with a compound which is an antagonist of the NMDA receptor.

U.S. Pat. No. 4,902,695 discloses certain competitive NMDA antagonists that are useful for the treatment of neurological disorders, including epilepsy, stroke, anxiety, cerebral ischemia, muscular spasms, and neurodegenerative diseases such as Alzheimer's disease and Huntington's disease.

U.S. Pat No. 5,192,751 discloses a method of treating urinary incontinence in a mammal which comprises administering an effective amount of a competitive NMDA antagonist.

## SUMMARY OF THE INVENTION

The invention relates BCAT inhibitor compounds of Formula I



I

- 3 -

wherein:

R<sub>3</sub> is H, or F, Br, alkyl, carboxy, alkoxy, substituted alkoxy,

R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, and R<sub>5</sub> are independently, H, halogen, alkyl, substituted alkyl, alkoxy,

substituted alkoxy, cyano, nitro, amino, alkylamine, and thioalkyl, Ar is

bicyclic heteroaryl, tricyclic heteroaryl, substituted bicyclic heteroaryl or

substituted tricyclic heteroaryl, except (6-methoxybenzofuran)-2-nyl, 3-

quinolinyl; or a pharmaceutically acceptable salt, ester, prodrug, or amide

thereof;

Ar is bicyclic heteroaryl, tricyclic heteroaryl, substituted bicyclic heteroaryl,

substituted tricyclic heteroaryl, except (6-methoxybenzofuran)-2-nyl or

3-quinolinyl;

where there is more than one stereoisomer, each chiral center may be independently R

or S; or a pharmaceutically acceptable salt, ester, prodrug, or amide thereof.

The invention also relates to compounds of Formula I, wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>,

and R<sub>5</sub> are hydrogen; R<sub>2</sub> or R<sub>4</sub> are alkyl; R<sub>2</sub> and R<sub>4</sub> are halogen; R<sub>1</sub> is alkyl and R<sub>2</sub> is

halogen; R<sub>5</sub> is halogen; and Ar is a tricyclic heteroaryl, bicyclic heteroaryl,

substituted bicyclic heteroaryl and substituted tricyclic heteroaryl.

The invention also relates to compounds of Formula I, wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>,

and R<sub>5</sub> are hydrogen, and Ar is a bicyclic heteroaryl; R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub> are

hydrogen, and Ar is a tricyclic heteroaryl; one of R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub> are halogen,

and Ar is a tricyclic heteroaryl; and two of R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub> are halogen, and Ar

is a tricyclic heteroaryl.

The invention also relates to compounds selected from:

Dibenzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2,3,4,9-Tetrahydro-1H-carbazole-6-carboxylic acid, 2-(phenylsulfonyl)-

hydrazide;

Benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

7-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

7-Ethoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

- 4 -

- 7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
Benzo[1,3] dioxole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
5 1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
1-Dibenzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;  
Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(2-methylphenyl)sulfonyl]hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;  
10 2-Dibenofurancarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]hydrazide;  
Dibenzofuran-2-carboxylic acid, 2-(3,5-dichlorophenylsulfonyl)hydrazide;  
Dibenzofuran-2-carboxylic acid, 2-(3-chloro-2-methyl-phenylsulfonyl)-  
hydrazide;  
15 Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
Quinoline-8-carboxylic acid, 2-(phenylsulfonyl)hydrazide;  
2-Benzofurancarboxylic acid, 5-chloro-2-[(4-fluorophenyl)]sulfonyl hydrazide;  
2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-fluoro-phenyl)sulfonyl]-  
hydrazide;  
20 2-Dibenzofurancarboxylic acid, 2-[(4-fluorophenyl)sulfonyl]hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(2-chloro-4-fluorophenyl)sulfonyl]-  
hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(3-cyanophenyl)sulfonyl]hydrazide;  
2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-cyanophenyl)sulfonyl]-  
25 hydrazide;  
2-Dibenzofurancarboxylic acid, 2-[(2,6-dichlorophenyl) sulfonyl]hydrazide;  
2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)hydrazide;  
2-Benzofurancarboxylic acid, 5-chloro-, 2-[(2-chloro-4-fluorophenyl)-  
sulfonyl]hydrazide;  
30 2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3,5-dichlorophenyl)-  
sulfonyl]hydrazide;

- 5 -

2-Dibenzofurancarboxylic acid, 2-[[3,5-bis(trifluoromethyl)phenyl]-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro, 2-[(3-chloro-4-fluorophenyl)-  
sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(5-fluoro-2-methylphenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3,4-difluorophenyl)sulfonyl]-hydrazide;

10 2-Dibenzofurancarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-fluorophenyl)sulfonyl]-hydrazide;

15 2-Benzofurancarboxylic acid, 5-chloro-2-[(3,4-difluorophenyl)sulfonyl]-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-chlorophenyl)sulfonyl]-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chlorophenyl)sulfonyl]-  
hydrazide;

20 2-Benzofurancarboxylic acid, 5-chloro-2-[(2-bromophenyl)sulfonyl]-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-bromophenyl)sulfonyl]-  
hydrazide;

25 2-Benzofurancarboxylic acid, 5-chloro-2-[(2-methylphenyl)sulfonyl]-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methylphenyl)sulfonyl]-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methoxyphenyl)sulfonyl]-  
hydrazide;

30 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chloro-2-methylphenyl)-  
sulfonyl]hydrazide;

- 6 -

2-Benzofurancarboxylic acid, 5-chloro-2-[(5-fluoro-2-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethoxy)phenyl]-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

10 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 6-methoxy-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl)hydrazide;

15 2-Benzofurancarboxylic acid, 3,5-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-,

2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(4-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

25 2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-, 2-(phenylsulfonyl)-hydrazide;

30 Naptho[2,3-*b*]furan-2-carboxylic acid, 3-methoxy-, 2-(phenylsulfonyl)-hydrazide;

- 7 -

2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl)sulfonyl hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

5           2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-methylphenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-methylphenyl) sulfonyl]-hydrazide;

10           2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-bromophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-bromophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

15           2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methoxy, 2-(phenylsulfonyl)-hydrazide;

20           2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

25           2-Benzofurancarboxylic acid, 5-methyl-, 2-[2-(methylphenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chloro-2-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[3-(bromophenyl)sulfonyl]-hydrazide;

30           2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

- 8 -

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)hydrazide;

5 2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methyl, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide;

10 2-Dibenzofurancarboxylic acid, 2-[(2-fluorophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-(2,6-difluorophenylsulfonyl)hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-bromophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide;

15 2-Dibenzofurancarboxylic acid, 2-[(4-methyl-3-nitrophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[2-chloro-5-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

20 2-Dibenzofurancarboxylic acid, 2-[(4-bromo-2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-carboxyphenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4,5-dichloro-2-thienyl)sulfonyl]-hydrazide;

25 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

30 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)-sulfonyl]hydrazide;

- 9 -

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)-sulfonyl]hydrazide;

10 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4,5-dibromo-2-thienyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]hydrazide;

15 2-Naphthalenecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[[4-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

20 2-Naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide;

25 2-Naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(4-pentylphenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[[2(trifluoromethyl)phenyl]sulfonyl]-hydrazide;

30 3-Quinolinecarboxylic acid, 2-[[2,5-bis(2,2,2-trifluoroethoxy)phenyl]-sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;

- 10 -

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]-sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;

5 3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide;

10 6-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide;

15 3-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

20 6-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide;

Benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

7-Ethoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

25 2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Isoquinoline-1-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1*H*-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-3-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

30 Quinoline-4-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1*H*-Indole-5-carboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide;

- 11 -

1*H*-Indole-5-carboxylic acid, 2-[(2-chlorophenyl)sulfonyl]hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(3-trifluoromethylphenyl)sulfonyl]-hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;  
 5     1*H*-Indole-5-carboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]-  
 hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(3trifluoromethoxyphenyl)sulfonyl]-  
 hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide;  
 10     1*H*-Indole-5-carboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide;  
 1*H*-Indole-5-carboxylic acid, 2-[(2-methylphenyl)sulfonyl]hydrazide;  
 6-Quinolinecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]hydrazide;  
 6-Quinolinecarboxylic acid, 2-[(3-trifluoromethylphenyl)sulfonyl]-hydrazide;  
 6-Quinolinecarboxylic acid, 2-[(3-trifluoromethoxyphenyl)sulfonyl]-hydrazide;  
 15     6-Quinolinecarboxylic acid, 2-[(2-methylphenyl)sulfonyl]hydrazide;  
 Dibenzofuran-2-carboxylic acid, 2-(4-fluoro-3-methylphenylsulfonyl)-  
 hydrazide;  
 7-Methoxy-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
 7-Chloro-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
 20     5-Methoxy-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
 5-Chloro-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
 5-Chloro-1*H*-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
 phenylsulfonyl]hydrazide;  
 5-Methyl-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
 25     5-Methyl-1*H*-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
 phenylsulfonyl]hydrazide; and  
 2-Naphthalenecarboxylic acid, 2-(phenylsulfonyl)hydrazide.

The invention also relates to a method of treating or preventing neuronal loss  
 associated with stroke, ischemia, CNS trauma, hypoglycemia and surgery, as well as  
 30     treating neurodegenerative diseases including Alzheimer's disease, amyotrophic lateral  
 sclerosis, Huntington's disease, Parkinson's disease and Down's syndrome, treating or

- 12 -

preventing the adverse consequences of the overstimulation of the excitatory amino acids, treating anxiety, psychosis, convulsions, chronic pain, neuropathic pain, diabetic retinopathy, glaucoma, CMV retinitis, urinary incontinence, and inducing anesthesia, as well as enhancing cognition, and preventing opiate tolerance and withdrawal symptoms, comprising administering to an animal in need of such treatment an effective amount of any one of the BCAT inhibitors of the present invention, or a pharmaceutically acceptable salt thereof.

The present invention also includes a pharmaceutical composition comprising a therapeutically effective amount of one or more compounds of Formula I and a pharmaceutically acceptable carrier.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention is predicated on the discovery that intraocular glutamate injection causes ganglion cell loss which is related to the early stages of diabetic retinopathy (Vorwerk et al., *IOVS*, 1996;37:1618-1624 and that an inhibitor of the branched chain amino acid-dependent aminotransferase pathway, specifically gabapentin, is effective in inhibiting the synthesis of glutamate (see example below), thus preventing diabetic retinopathy. Accordingly, the present invention provides a method for the prophylactic and therapeutic treatment of diabetic retinopathy, including treatment at the pre-diabetic retinopathy stage, the nonproliferative diabetic retinopathy stage, and the proliferative diabetic retinopathy stage. By "prophylactic" is meant the protection, in whole or in part, against diabetic retinopathy, in particular diabetic macular edema. By "therapeutic" is meant the amelioration of diabetic retinopathy, itself, and the protection, in whole or in part, against further diabetic retinopathy, in particular diabetic macular edema.

The method comprises the administration of an inhibitor of the branched chain amino acid-dependent aminotransferase pathway in an amount sufficient to treat the neurodegenerative disease or condition, for example, to treat the retina for retinopathy prophylactically or therapeutically. Any inhibitor of the branched chain amino acid-dependent aminotransferase pathway can be used in the method of the present

- 13 -

invention as long as it is safe and efficacious. Herein, "branch chain amino acid-dependent aminotransferase (BCAT) inhibitor" will be used to refer to such compounds and is intended to encompass all compounds that affect the branch chain amino acid-dependent aminotransferase pathway at any and all points in the pathway.

5            Preferably, the BCAT inhibitor is a compound of Formula I as described above, or a pharmaceutically acceptable, BCAT pathway-inhibiting analogue or prodrug thereof or a pharmaceutically acceptable salt, ester, or amide of any of the foregoing.

10           Unless otherwise expressly stated, the following definitions are adhered to throughout this disclosure.

             "Alkyl" means a straight or branched hydrocarbon radical having from 1 to 10 carbon atoms (unless stated otherwise) and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, iso-pentyl, n-hexyl, and the like. "Halogen" includes fluoro, chloro, bromo, and iodo.

15           "Alkenyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and one double bond and includes ethenyl, 3-buten-1-yl, 2-ethenylbutyl, 3-hexen-1-yl, and the like.

             "Alkynyl" means straight and branched hydrocarbon radicals having from 2 to 6 carbon atoms and one triple bond and includes ethynyl, 3-butyn-1-yl, propynyl, 20           2-butyn-1-yl, 3-pentyn-1-yl, and the like.

             "Cycloalkyl" means a monocyclic or polycyclic hydrocarbyl group such as cyclopropyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclobutyl, adamantyl, norpinanyl, decalanyl, norbornyl, cyclohexyl, and cyclopentyl. Such groups can be substituted with groups such as hydroxy, keto, alkoxy, hydroxyalkyl and the like. Also included are 25           rings in which 1 to 3 heteroatoms replace carbons. Such groups are termed "heterocyclyl," which means a cycloalkyl group also bearing at least one heteroatom selected from O, S, or NR<sub>2</sub>, wherein R<sub>2</sub> is as defined above with respect to substituents of the phenyl group of Formula I. Suitable examples include substituted or unsubstituted oxiranyl, pyrrolidinyl, piperidyl, tetrahydropyran, piperazinyl, 30           acylpiperazinyl, pyrrolidinyl, and morpholine.

- 14 -

“Alkoxy” refers to the alkyl groups mentioned above bound through oxygen, examples of which include methoxy, ethoxy, isopropoxy, tert-butoxy, and the like. In addition, alkoxy refers to polyethers such as  $-O-(CH_2)_2-O-CH_3$ , and the like.

“Alkanoyl” groups are alkyl linked through a carbonyl, ie,  $C_1-C_5-C(O)-$ . Such groups include formyl, acetyl, propionyl, butyryl, and isobutyryl.

“Acyl” means an alkyl or aryl (Ar) group bonded through a carbonyl group, ie,  $R-C(O)-$ . For example, acyl includes a  $C_1-C_6$  alkanoyl, including substituted alkanoyl, wherein the alkyl portion can be substituted by  $NR_4R_5$  or a carboxylic or heterocyclic group.  $R_4$  and  $R_5$  are as defined above with respect to substituents of the phenyl group of Formula I. Typical acyl groups include acetyl, benzoyl, and the like.

The alkyl, alkenyl, alkoxy, and alkynyl groups described above are optionally substituted, preferably by 1 to 3 groups selected from  $NR_4R_5$ , phenyl, substituted phenyl (wherein the substituent on the phenyl group may be selected from 1 to 3 groups selected from  $NR_4R_5$ , phenyl, substituted phenyl, thio  $C_1-C_6$  alkyl,  $C_1-C_6$  alkoxy, hydroxy, carboxy,  $C_1-C_6$  alkoxycarbonyl, halo, nitrile, cycloalkyl, and a 5- or 6-membered carbocyclic ring or heterocyclic ring having 1 or 2 heteroatoms selected from nitrogen, substituted nitrogen, oxygen, and sulfur), thio  $C_1-C_6$  alkyl,  $C_1-C_6$  alkoxy, hydroxy, carboxy,  $C_1-C_6$  alkoxycarbonyl, halo, nitrile, cycloalkyl, and a 5- or 6-membered carbocyclic ring or heterocyclic ring having 1 or 2 heteroatoms selected from nitrogen, substituted nitrogen, oxygen, and sulfur.  $R_4$  and  $R_5$  are as defined above with respect to substituents of the phenyl group of Formula I. “Substituted nitrogen” means nitrogen bearing  $C_1-C_6$  alkyl or  $(CH_2)_nPh$  where n is 1, 2, or 3. Perhalo and polyhalo substitution is also embraced.

Examples of substituted alkyl groups include 2-aminoethyl, pentachloroethyl, trifluoromethyl, 2-diethylaminoethyl, 2-dimethylaminopropyl, ethoxycarbonylmethyl, 3-phenylbutyl, methanysulfanylmethyl, methoxymethyl, 3-hydroxypentyl, 2-carboxybutyl, 4-chlorobutyl, 3-cyclopropylpropyl, pentafluoroethyl, 3-morpholinopropyl, piperazinylmethyl, and 2-(4-methylpiperazinyl)ethyl.

- 15 -


Examples of substituted alkynyl groups include 2-methoxyethynyl, 2-ethylsulfanyethynyl, 4-(1-piperazinyl)-3-(butynyl), 3-phenyl-5-hexynyl, 3-diethylamino-3-butynyl, 4-chloro-3-butynyl, 4-cyclobutyl-4-hexenyl, and the like.

Typical substituted alkoxy groups include aminomethoxy, trifluoromethoxy, 2-diethylaminoethoxy, 2-ethoxycarbonylethoxy, 3-hydroxypropoxy, 6-carboxhexyloxy, and the like.


Further, examples of substituted alkyl, alkenyl, and alkynyl groups include dimethylaminomethyl, carboxymethyl, 4-dimethylamino-3-buten-1-yl, 5-ethylmethylamino-3-pentyn-1-yl, 4-morpholinobutyl, 4-tetrahydropyridinylbutyl, 3-imidazolidin-1-ylpropyl, 4-tetrahydrothiazol-3-yl-butyl, phenylmethyl, 3-chlorophenylmethyl, and the like.

The terms "Ar" and "aryl" refer to unsubstituted and substituted aromatic groups. Heteroaryl groups have from 4 to 9 ring atoms, from 1 to 4 of which are independently selected from the group consisting of O, S, and N. Preferred heteroaryl groups have 1 or 2 heteroatoms in a 5- or 6-membered aromatic ring. Mono and bicyclic aromatic ring systems are included in the definition of aryl and heteroaryl. Typical aryl and heteroaryl groups include phenyl, 3-chlorophenyl, 2,6-dibromophenyl, pyridyl, 3-methylpyridyl, benzothienyl, 2,4,6-tribromophenyl, 4-ethylbenzothienyl, furanyl, 3,4-diethylfuranyl, naphthyl, 4,7-dichloronaphthyl, pyrrole, pyrazole, imidazole, thiazole, and the like.

Examples are Ar include but are not limited to:



- 16 -



wherein R<sub>6</sub> is alkoxy, such as methoxy or ethoxy, substituted alkoxy, halogen, such as chloro, aryloxy, substituted aryloxy, alkylamino, substituted alkylamino, arylalkylamino, substituted arylalkylamino, amide or substituted amide where in the substituents may be as defined above.

Preferred Ar groups are phenyl and phenyl substituted by 1, 2, or 3 groups independently selected from the group consisting of alkyl, alkoxy, thio, thioalkyl, hydroxy, -COOR<sub>7</sub>, amino of the formula -NR<sub>4</sub>R<sub>5</sub>, and T(CH<sub>2</sub>)<sub>m</sub>QR<sub>4</sub> or T(CH<sub>2</sub>)<sub>m</sub>CO<sub>2</sub>R<sub>4</sub> wherein m is 1 to 6, T is O, S, NR<sub>4</sub>, N(O)R<sub>4</sub>, or CR<sub>4</sub>R<sub>5</sub>, Q is O, S, NR<sub>5</sub>, N(O)R<sub>5</sub>, wherein R<sub>4</sub> and R<sub>5</sub> are as described above, and R<sub>7</sub> is alkyl or substituted alkyl, for example, methyl, trichloroethyl, diphenylmethyl, and the like. The alkyl and alkoxy groups can be substituted as defined above. For example, typical groups are carboxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, hydroxyalkoxy, and alkoxyalkyl.

The symbol “-” means a bond.

The term “patient” means all animals including humans. Examples of patients include humans, cows, dogs, cats, goats, sheep, and pigs.

The compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds of the present invention possess one or more chiral centers and each center may exist in the R(D) or S(L) configuration. The present invention includes all enantiomeric and epimeric forms as well as the appropriate mixtures thereof.

The compounds of Formula I are capable of further forming both pharmaceutically acceptable formulations comprising salts, esters, amides, and

- 17 -

prodrugs. As used herein, the term “pharmaceutically acceptable salts, esters, amides, and prodrugs” refers to those carboxylate salts, amino acid addition salts, esters, amides, and prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term “salts” refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed and including, but not limited to, acid addition and/or base salts, solvents and N-oxides of a compound of Formula I. This invention also provides pharmaceutical formulations comprising a compound of Formula I together with a pharmaceutically acceptable carrier, diluent, or excipient therefor. All of these forms are within the present invention.

Pharmaceutically acceptable acid addition salts of the compounds of Formula I include salts derived from inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, phosphorus, and the like, as well as the salts derived from organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Also contemplated are the salts of amino acids such as arginate, gluconate, galacturonate, and the like; see, for example, Berge, et al., “Pharmaceutical Salts,” *J. of Pharmaceutical Science*, 1977;66:1-19.

- 18 -

The acid addition salts of the basic compounds are prepared by contacting the free base form with a sufficient amount of the desired acid to produce the salt in the conventional manner. The free base form may be regenerated by contacting the salt form with a base and isolating the free base in the conventional manner. The free base forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free base for purposes of the present invention.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metal hydroxides, or of organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, N-methylglucamine, and procaine; see, for example, Berge et al., supra., 1977.

The base addition salts of acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in a conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention.

Examples of pharmaceutically acceptable, non-toxic esters of the compounds of this invention include C<sub>1</sub>-C<sub>6</sub>alkyl esters wherein the alkyl group is a straight or branched chain. Acceptable esters also include C<sub>5</sub>-C<sub>7</sub>cycloalkyl esters as well as arylalkyl esters such as, but not limited to benzyl. C<sub>1</sub>-C<sub>4</sub>alkyl esters are preferred. Esters of the compounds of the present invention may be prepared according to conventional methods.

Examples of pharmaceutically acceptable, non-toxic amides of the compounds of this invention include amides derived from ammonia, primary C<sub>1</sub>-C<sub>6</sub>alkyl amines and secondary C<sub>1</sub>-C<sub>6</sub>dialkyl amines wherein the alkyl groups are straight or branched chain. In the case of secondary amines the amine may also be in the form of a 5- or 6-membered heterocycle containing one nitrogen atom. Amides derived from

- 19 -

ammonia, C<sub>1</sub>-C<sub>3</sub>alkyl primary amines, and C<sub>1</sub>-C<sub>2</sub>dialkyl secondary amines are preferred. Amides of the compounds of the invention may be prepared according to conventional methods.

The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above Formulae, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference. In general, a prodrug is a drug which has been chemically modified and may be biologically inactive at its site of action, but which may be degraded or modified by one or more enzymatic or other *in vivo* processes to the parent bioactive form.

A therapeutically effective amount is an amount of a compound of Formula I that when administered to a patient, ameliorates a symptom of the disease.

The BCAT inhibitor, which is preferably a compound of Formula I, a BCAT pathway-inhibiting analogue of Formula I, a BCAT pathway-inhibiting prodrug of Formula I, or a pharmaceutically acceptable salt of any of the foregoing, can be administered in accordance with the present inventive method by any suitable route. Suitable routes of administration include systemic, such as orally or by injection, topical, intraocular, periocular (e.g., subTenon's), subconjunctival, subretinal, suprachoroidal and retrobulbar. The manner in which the BCAT inhibitor is administered is dependent, in part, upon whether the treatment of retinopathy is prophylactic or therapeutic. The manner in which the BCAT inhibitor is administered for therapeutic treatment of retinopathy is dependent, in part, upon the cause of the retinopathy.

For example, given that diabetes is the leading cause of retinopathy, the BCAT inhibitor can be administered prophylactically as soon as the pre-diabetic retinopathy state is detected. For the prophylactic treatment of retinopathy that can result from diabetes, the BCAT inhibitor is preferably administered systemically, e.g., orally or by injection. For the therapeutic treatment of nonproliferative diabetic retinopathy, the

- 20 -

BCAT inhibitor can be administered systemically, e.g., orally or by injection, or intraocularly. Proliferative diabetic retinopathy can be therapeutically treated by the administration of the BCAT inhibitor intraocularly, topically, subconjunctivally or periocularly (e.g., subTenon's), for example. The BCAT inhibitor is preferably administered intraocularly, topically, subconjunctivally or periocularly (e.g., subTenon's) for the prophylactic or therapeutic treatment of retinopathy before, during or after surgical removal from an eye of scar tissue generated during neovascularization during the proliferative diabetic stage.

The BCAT inhibitor is preferably administered as soon as possible after it has been determined that an animal, such as a mammal, specifically a human, is at risk for retinopathy (prophylactic treatment) or has begun to develop retinopathy (therapeutic treatment). Treatment will depend, in part, upon the particular BCAT inhibitor used, the amount of the BCAT inhibitor administered, the route of administration, and the cause and extent, if any, of retinopathy realized.

One skilled in the art will appreciate that suitable methods of administering a BCAT inhibitor, which is useful in the present inventive method, are available. Although more than one route can be used to administer a particular BCAT inhibitor, a particular route can provide a more immediate and more effective reaction than another route. Accordingly, the described routes of administration are merely exemplary and are in no way limiting.

The dose administered to an animal, particularly a human, in accordance with the present invention should be sufficient to effect the desired response in the animal over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors, including the strength of the particular BCAT inhibitor employed, the age, species, condition or disease state, and body weight of the animal, as well as the amount of the retina about to be affected or actually affected by retinopathy. The size of the dose also will be determined by the route, timing and frequency of administration as well as the existence, nature, and extent of any adverse side effects that might accompany the administration of a particular BCAT inhibitor and the desired physiological effect. It will be appreciated by one of ordinary skill in

- 21 -

the art that various conditions or disease states, in particular, chronic conditions or disease states, may require prolonged treatment involving multiple administrations.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. The present inventive method will typically involve the administration of from about 1 mg/kg/day to about 100 mg/kg/day, preferably from about 15 mg/kg/day to about 50 mg/kg/day, if administered systemically. Intraocular administration typically will involve the administration of from about 0.1 mg total to about 5 mg total, preferably from about 0.5 mg total to about 1 mg total. A preferred concentration for topical administration is 100  $\mu$ M.

Compositions for use in the present inventive method preferably comprise a pharmaceutically acceptable carrier and an amount of a BCAT inhibitor sufficient to treat retinopathy prophylactically or therapeutically. The carrier can be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration. It will be appreciated by one of ordinary skill in the art that, in addition to the following described pharmaceutical compositions, the BCAT inhibitor can be formulated as polymeric compositions, inclusion complexes, such as cyclodextrin inclusion complexes, liposomes, microspheres, microcapsules and the like (see, e.g., U.S. Pat. Nos. 4,997,652; 5,185,152; and 5,718,922).

The BCAT inhibitor can be formulated as a pharmaceutically acceptable acid addition salt. Examples of pharmaceutically acceptable acid addition salts for use in the pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, and arylsulphonic, for example p-toluenesulphonic, acids.

The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically

- 22 -

acceptable carrier be one which is chemically inert to the BCAT inhibitor and one which has no detrimental side effects or toxicity under the conditions of use.

The choice of excipient will be determined in part by the particular BCAT inhibitor, as well as by the particular method used to administer the composition.

5 Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention. The following formulations are merely exemplary and are in no way limiting.

Injectable formulations are among those that are preferred in accordance with the present inventive method. The requirements for effective pharmaceutically carriers  
10 for injectable compositions are well-known to those of ordinary skill in the art (see *Pharmaceutics and Pharmacy Practice*, J. B. Lippincott Co., Philadelphia, PA, Banker and Chalmers, eds., pages 238-250 (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986). It is preferred that such injectable compositions be administered intramuscularly, intravenously, or intraperitoneally.

15 Topical formulations are well-known to those of skill in the art. Such formulations are suitable in the context of the present invention for application to the skin. The use of patches, corneal shields (see, e.g., U.S. Pat. No. 5,185,152), and ophthalmic solutions (see, e.g., U.S. Pat. No. 5,710,182) and ointments, e.g., eye drops, is also within the skill in the art.

20 Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid  
25 formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent.

Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium  
30 phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginate acid, microcrystalline cellulose, acacia,

- 23 -

gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can  
5 comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

The effectiveness of an orally administered drug is dependent upon the drug's  
10 efficient transport across the mucosal epithelium and its stability in entero-hepatic circulation. Drugs that are effective after parenteral administration but less effective orally, or whose plasma half-life is considered too short, may be chemically modified into a prodrug form.

Formulations suitable for parenteral administration include aqueous and non-  
15 aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The inhibitor can be administered in a physiologically acceptable diluent in a  
20 pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, dimethylsulfoxide, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid  
25 ester or glyceride, or an acetylated fatty acid glyceride, with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants. Oils, which can be used in parenteral formulations include petroleum, animal,  
30 vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral.

- 24 -

Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters.

Suitable soaps for use in parenteral formulations include fatty alkali metals, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl-p-aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations will typically contain from about 0.5 to about 25% by weight of the active ingredient in solution. Preservatives and buffers may be used. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17.

The quantity of surfactant in such formulations will typically range from about 5 to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.


Such compositions can be formulated as intraocular formulations, sustained-release formulations or devices (see, e.g., U.S. Pat. No. 5,378,475). For example, gelatin, chondroitin sulfate, a polyphosphoester, such as bis-2-hydroxyethyl-terephthalate (BHET), or a polylactic-glycolic acid (in various proportions) can be

- 25 -

used to formulate sustained-release formulations. Implants (see, e.g., U.S. Pat. Nos. 5,443,505; 4,853,224; and 4,997,652), devices (see, e.g., U.S. Pat. Nos. 5,554,187; 4,863,457; 5,098,443; and 5,725,493), such as an implantable device, e.g., a mechanical reservoir, an intraocular device or an extraocular device with an intraocular conduit (e.g., 100  $\mu$  - 1 mm in diameter), or an implant or a device comprised of a polymeric composition as described above, can be used.

The present inventive method also can involve the co-administration of other pharmaceutically active compounds. By "co-administration" is meant administration before, concurrently with, e.g., in combination with the BCAT inhibitor in the same formulation or in separate formulations, or after administration of a BCAT inhibitor as described above. For example, corticosteroids, e.g., prednisone, methylprednisolone, dexamethasone, or triamcinalone acetonide, or noncorticosteroid anti-inflammatory compounds, such as ibuprofen or flubiprofen, can be co-administered. Similarly, vitamins and minerals, e.g., zinc, antioxidants, e.g., carotenoids (such as a xanthophyll carotenoid like zeaxanthin or lutein), and micronutrients can be co-administered.

Compounds of the present invention may be obtained by reaction between a hydrazide of the formula  $\text{ArC(O)NH}_2\text{NH}_2$  with a sulfonyl halide of the formula:



Where R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and Ar are as defined above and Hal is halogen, preferably chlorine. The hydrazide starting materials may be prepared in conventional manner by reaction of the corresponding carboxylic acid with hydrazine. The reaction between the hydrazide and the sulfonyl halide may be carried out in manner known for reaction between these classes of compounds and may conveniently be carried out in


- 26 -

an organic solvent and where desired using a catalyst such as an amine catalyst, for example dimethylaminopyridine or N,N-diisopropylethylamine. Such a sequence of reactions is illustrated by the following reaction scheme:

5

General synthetic scheme for preparing compounds of Formula I


Scheme 1



## EXAMPLE 1

## 10 Dibenzo-furan-2-carboxylic acid 2-(phenylsulfonyl)hydrazide


Step 1. Dibenzo-furan-2-carboxylic acid (**1a**) was prepared in accordance with the methods of Ames et al., *Synthesis*, 1983:234.



MS: 212.0 (M+1 for C<sub>13</sub>H<sub>8</sub>O<sub>3</sub>); mp: 248-249°C; IR (KBr, cm<sup>-1</sup>): 2825, 1668, 1434,

15 1310. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.39-7.43 (m, 1H), 7.51-7.56 (m, 1H), 7.69-7.77 (m, 2H), 8.06-8.10 (m, 1H), 8.26 (d, 1H, J = 7.6 Hz), 8.74 (d, 1H, J = 1.7 Hz).

Step 2. Dibenzo-furan-2-carboxylic acid hydrazide (**1b**)



Synthesis of dibenzofuran-2-carboxylic acid hydrazide: Dibenzo-furan-2-carboxylic acid (3.50 g, 16.5 mmol) was dissolved in DMF (160 mL), treated with N-methyl-morpholine (4.0 mL, 36.3 mmol), cooled to 0°C, treated with isobutyl

20


- 27 -

chloroformate (2.35 mL, 18.1 mmol), and stirred for 10 minutes. Hydrazine (10.4 mL, 330.0 mmol) was added, and the reaction was allowed to warm to room temperature and stir for 3 hours. The reaction was then diluted with EtOAc (400 mL), washed with saturated sodium bicarbonate solution and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo to give 3.40 g (91%) of the desired product as a white solid which was used without further purification.

MS: 227.0 (M+1 for C<sub>13</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub>); mp: 204-205°C; TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.34 (8% MeOH/CH<sub>2</sub>Cl<sub>2</sub>); HPLC (C18 column, 1:1 CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% TFA) 94.76%,

RT = 2.196 min. HRMS (calc): 227.0820 (found): 227.0827. IR (KBr, cm<sup>-1</sup>): 3303, 1620, 1542, 1472, 1192. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.45 (br, 2H), 7.39-7.43 (m, 1H), 7.51-7.55 (m, 1H), 7.72 (t, 2H, J = 8.5 Hz), 7.96 (dd, 1H, J = 8.5, 1.7 Hz), 8.15 (d, 1H, J = 8.3 Hz), 8.61 (d, 1H, J = 1.7 Hz), 9.83 (s, 1H).


Step 3. Dibenzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



Dibenzofuran-2-carboxylic acid hydrazide (**1b**, 0.3 g, 1.33 mmol) was dissolved in DMF (9 mL) and treated with Hunig's base (0.93 mL, 5.32 mmol). The reaction was cooled to 0°C, treated with benzenesulfonyl chloride (0.186 mL, 1.46 mmol) and a catalytic amount of dimethylaminopyridine (5 mg), then allowed to warm to room temperature and stir overnight. The reaction was then diluted with EtOAc (125 mL), washed with saturated sodium bicarbonate solution and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo. The crude material was chromatographed on silica gel eluting with 6% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 0.25 g (50%) of the desired product. MS: 367.0 (M+1 for C<sub>19</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 216-218°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.58 (1:1 hexane/EtOAc); Analysis (C<sub>19</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S) (calc) C: 62.29, H: 3.85, N: 7.65; (found) C: 62.16, H: 3.88, N: 7.60. IR (KBr, cm<sup>-1</sup>): 3334, 3157, 1659, 1414, 1203, 1172. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.41 (t, 1H, J = 7.6 Hz), 7.49-7.61 (m, 4H), 7.71 (t, 2H, J = 8.5 Hz), 7.82-7.83 (m, 3H), 8.16 (d, 1H, J = 7.8 Hz), 8.49 (s, 1H).

- 28 -

## EXAMPLE 2


2,3,4,9-Tetrahydro-1*H*-carbazole-6-carboxylic acid 2-(phenylsulfonyl)hydrazideSynthesis of 2,3,4,9-tetrahydro-1*H*-carbazole-6-carboxylic acid

2-(phenylsulfonyl)hydrazide: Example 2 was synthesized in accordance with the methods of Example 1 except that 6,7,8,9-tetrahydro-5*H*-carbazole-3-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide (56% yield). MS: 370.1 (M+1 for C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub>S<sub>1</sub>); mp: >250°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.28 (1:1 hexane/EtOAc); HPLC (C18 column, 1:1 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 99.25%,

RT = 7.748 min. IR (KBr, cm<sup>-1</sup>): 3403, 3340, 3124, 2941, 1657, 1422, 1165. Analysis (C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub>S<sub>1</sub>); (calc) C: 61.77, H: 5.18, N: 11.37; (found) C: 61.56, H: 5.15, N: 11.17. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.76-1.80 (m, 4H), 2.46-2.65 (m, 4H), 7.17 (d, 1H, J = 8.3 Hz), 7.33 (d, 1H, J = 8.5 Hz), 7.44-7.48 (m, 2H), 7.56 (t, 1H, J = 7.1 Hz), 7.76-7.81 (m, 3H), 9.80 (s, 1H), 10.38 (s, 1H), 10.91 (s, 1H).

## EXAMPLE 3

Benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



Benzofuran-2-carboxylic acid hydrazide (0.4 g, 2.27 mmol) was suspended in THF (25 mL). N,N-diisopropylethyl amine (1.58 mL, 9.08 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.32 mL, 2.50 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was orange after it stirred for 1 hour, and DMF (20 mL) was added and the solution became clear. After 4 hours of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified

- 29 -

by flash chromatography eluting with 3:1 hexane/EtOAc. The desired product was isolated (0.38 g, 53.0%).

MS: 317.0 (M+1 for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.23 (1:1 hexane/EtOAc). mp:

175-180°C; HRMS: 317.0598 (M+1 for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S), HPLC: (30% H<sub>2</sub>O/70%


CH<sub>3</sub>CN/0.1% TFA), Ret. Time 2.911, Purity: 93.12%. IR (KBr, cm<sup>-1</sup>) 3338, 3131,

1676, 1581, 1349, 1163. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.23-7.67 (m, 9H), 7.93 (d;

2H, J = 5.6 Hz), 8.52 (s, 1H).

#### EXAMPLE 4

5-Methoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)hydrazide



Synthesis of 5-methoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-hydrazide: Example 4 was synthesized in accordance with the methods of Example 1 except that 5-methoxy-benzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid hydrazide (3.2% yield).

MS: 347.0 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S<sub>1</sub>); glassy solid; TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.16 (3:2 hexane/EtOAc); HPLC (C18 column, 1:1 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 99.12%,

RT = 5.813 min. HRMS (calc for M+1) 347.0701 (found) 347.0696. IR (KBr, cm<sup>-1</sup>):

3302, 3139, 2927, 1666, 1582, 1168. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.75 (s, 3H), 7.01 (dd,


1H, J = 9.0, 2.4 Hz), 7.21 (d, 1H, J = 2.4 Hz), 7.49-7.53 (m, 4H), 7.61 (t, 1H, J = 7.6

Hz), 7.79-7.81 (m, 2H), 10.13 (br, 1H), 10.85 (br, 1H).

- 30 -

## EXAMPLE 5

## 7-Methoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)hydrazide




Synthesis of 7-methoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-  
5 hydrazide: Example 5 was synthesized in accordance with the methods of Example 1  
except that 7-methoxy-benzofuran-2-carboxylic acid was used instead of  
dibenzofuran-4-carboxylic acid (64% yield).

MS: 347.0 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S<sub>1</sub>); mp: 164-165°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.35 (1:1  
hexane/EtOAc); HPLC (C18 column, 40:60 CH<sub>3</sub>CN/H<sub>2</sub>O+0.05% TFA) 99.87%, RT =  
10 11.355 min. HRMS (calc for M+1) 347.0701 (found) 347.0702. IR (KBr, cm<sup>-1</sup>): 3232,  
1675, 1588, 1490, 1166. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 4.03 (s, 3H), 6.94 (dd, 1H, J = 7.1, 2.0  
Hz), 7.21-7.26 (m, 2H), 7.34-7.37 (m, 2H), 7.45-7.50 (m, 2H), 7.58-7.63 (m, 1H),  
7.93-7.96 (m, 2H), 8.61 (d, 1H, J = 6.1 Hz).

## EXAMPLE 6

## 7-Ethoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)hydrazide



7-Ethoxy-benzofuran-2-carboxylic acid hydrazide (0.5 g, 2.27 mmol) was  
dissolved in THF (25 mL). N,N-diisopropylethylamine (1.58 mL, 9.08 mmol) was  
added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.32 mL,  
20 2.50 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was  
allowed to warm to room temperature. The solution was orange after it stirred for  
1 hour. After 3 hours of stirring, the solution was washed with saturated sodium  
bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated

- 31 -

down and purified by flash chromatography eluting with 1:1 hexane/EtOAc. The desired product was isolated (0.05 g, 5.6%).

MS: 361.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.17 (1:1 hexane/EtOAc); mp:

223-224°C. HRMS: (calc) 361.0558, (found) 361.0853 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S)


HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 2.596, Purity: 98.71%. IR

(KBr, cm<sup>-1</sup>) 3268, 3124, 2982, 1666, 1586, 1344, 1166, 1087, 724. <sup>1</sup>H NMR

(400 MHz, DMSO-d<sub>6</sub>) δ 1.37 (t, 3H, J = 6.8 Hz), 4.19 (q, 2H, J = 7.1 Hz), 7.01 (d, 1H, J = 8.0 Hz), 7.16-7.26 (m, 2H), 7.49-7.62 (m, 4H), 7.81 (d, 1H, J = 7.3 Hz), 10.17 (s, 1H), 10.92 (s, 1H).

## EXAMPLE 7

7-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



7-Chloro-benzofuran-2-carboxylic acid hydrazide (0.5 g, 2.38 mmol) was suspended in THF (26 mL). N,N-diisopropylethylamine (1.66 mL, 9.52 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.33 mL, 2.62 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was yellow after it stirred for 2 hours, and DMF (10 mL) was added. After another hour of stirring, the solution was diluted with EtOAc (150 mL), and washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3:1 hexane/EtOAc. The desired product was isolated (0.04 g, 4.6%).

MS: 351.0 (M+1 for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.57 (1:1 hexane/EtOAc); mp: 223-224°C; HRMS: (calc) 351.0206, (found) 351.0215 (M+1 for


C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl) HPLC: Ret. Time 9.862, Purity: 98.26%. IR (KBr, cm<sup>-1</sup>) 3291,

- 32 -

3033, 1670, 1344, 1164.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24-7.64 (m, 7H), 7.94 (d, 2H,  $J = 7.1$  Hz), 8.62 (s, 1H), 9.12 (s, 1H).

## EXAMPLE 8

## 5-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide




5-Chloro-benzofuran-2-carboxylic acid hydrazide (0.4 g, 1.90 mmol) was suspended in THF (21 mL). N,N-diisopropylethylamine (1.32 mL, 7.6 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.27 mL, 2.09 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was orange after it stirred for 2 hours, and DMF (10 mL) was added and the solution became clear. After 15 minutes of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over  $\text{Na}_2\text{SO}_4$ . The reaction was then concentrated down and purified by flash chromatography eluting with 2:1 hexane/EtOAc. The desired product was isolated (0.03 g, 3.8%). MS: 350.9 ( $\text{M}+1$  for  $\text{C}_{15}\text{H}_{11}\text{N}_2\text{O}_4\text{SCl}$ ). TLC:  $\text{SiO}_2$ ,  $R_f = 0.55$  (1:1 hexane/EtOAc); mp: 181-183°C; HRMS: (calc) 351.0206, (found) 351.0206 ( $\text{M}+1$  for  $\text{C}_{15}\text{H}_{11}\text{N}_2\text{O}_4\text{SCl}$ ) HPLC: (50%  $\text{H}_2\text{O}$ /50%  $\text{CH}_3\text{CN}$ /0.1% TFA) Ret. Time 8.755, Purity: 97.13%. IR (KBr,  $\text{cm}^{-1}$ ) 3291, 3066, 1670, 1344, 1164.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  4.79 (br, 1H), 7.24-7.69 (m, 7H), 7.92 (d, 2H,  $J = 7.6$  Hz), 8.51 (br, 1H).

- 33 -

## EXAMPLE 9

2,3-Dihydro-benzofuran-5-carboxylic acid 2-(phenylsulfonyl)hydrazide




2,3-Dihydro-benzofuran-5-carboxylic acid hydrazide (0.25 g, 1.40 mmol) was dissolved in THF (16 mL). N,N-diisopropylethylamine (0.98 mL, 5.6 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.20 mL, 1.54 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 2 hours of stirring, the solution was diluted with EtOAc (200 mL), and washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 1:1 hexane/EtOAc. The desired product was isolated (0.09 g, 20.0%).

MS: 319.0 (M+1 for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S). mp: 181-183 °C; TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.46 (1:1 hexane/EtOAc); HRMS: (calc) 319.0752, (found) 319.0761 (M+1 for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S).  
HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time 9.862, Purity: 95.04%. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 3.13 (t, 2H, J = 8.8 Hz), 4.54 (t, 2H, J = 8.8 Hz), 6.74 (d, 1H, J = 8.3 Hz), 7.46-7.57 (m, 5H), 7.77 (d, 2H, J = 7.0 Hz), 9.85 (s, 1H), 10.42 (s, 1H).

## EXAMPLE 10

Benzo[1,3] dioxole-5-carboxylic acid 2-(phenylsulfonyl)hydrazide



Benzo[1,3] dioxole-5-carboxylic acid hydrazide (0.5 g, 3.33 mmol) was suspended in THF (37 mL). N,N-diisopropylethylamine (2.32 mL, 13.32 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.47 mL,


- 34 -

3.66 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was orange after it stirred for 1 hour, and DMF (5 mL) was added and the solution became clear. After 2 hours of stirring, the solution was diluted with EtOAc (100 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3:1 hexane/EtOAc. The desired product was isolated (0.09 g, 8.8%).

MS: 321.0 (M+1 for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.59 (1:1 hexane/EtOAc); mp: 156-158°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 5.359, Purity: 95.27%. Analysis: (C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>S) (calc) C: 52.49, H: 3.78, N: 8.75 (found) C: 52.61, H: 4.06, N: 8.35. IR (KBr, cm<sup>-1</sup>) 3210, 2920, 1643, 1502, 1390, 1251, 1168, 1029. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 6.05 (s, 2H), 6.92 (d, 1H, J = 8.1 Hz), 7.15 (s, 1H), 7.24 (d, 1H, J = 8.1 Hz), 7.48 (t, 2H, J = 7.8 Hz), 7.58 (t, 1H, J = 7.6 Hz), 7.77 (d, 2H, J = 7.1 Hz), 9.92 (s, 1H), 10.49 (s, 1H).

## EXAMPLE 11

1*H*-Indole-5-carboxylic acid 2-(phenylsulfonyl)hydrazide




1*H*-Indole-5-carboxylic acid hydrazide (0.5 g, 2.85 mmol) was suspended in THF (32 mL). N,N-diisopropylethylamine (1.99 mL, 11.40 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.36 mL, 2.85 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature, and DMF (3 mL) was added. After 3 hours of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 2:1 hexane/EtOAc. The desired product was isolated (0.41 g, 23.8%).

- 35 -

MS: 316.0 (M+1 for C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.38 (1:1 hexane/EtOAc); mp: 209-211°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 4.102, Purity: 96.75%. Analysis (C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S) (calc) C: 57.13, H: 4.16, N: 13.32 (found) C: 57.28, H: 4.08, N: 13.06. IR (KBr, cm<sup>-1</sup>) 3293, 3168, 1649, 1312, 1168. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 6.48 (s, 1H), 7.33-7.41 (m, 3H), 7.47 (t, 2H, J = 7.3 Hz), 7.57 (t, 1H, J = 7.4 Hz), 7.80 (d, 2H, J = 7.3 Hz), 7.94 (s, 1H), 9.84 (s, 1H), 10.45 (s, 1H), 11.33 (s, 1H).

## EXAMPLE 12

1-Dibenzofurancarboxylic acid 2-(phenylsulfonyl)hydrazide




Synthesis of 1-dibenzofurancarboxylic acid 2-(phenylsulfonyl)hydrazide:

Example 12 was synthesized in accordance with the methods of Example 1 except that dibenzofuran-1-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide (8.4%).

MS: 367.0 (M+1 for C<sub>19</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>); glassy solid; TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.77 (8% MeOH/CH<sub>2</sub>Cl<sub>2</sub>); HPLC (C18 column, 7:3 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 91.94%, RT = 4.008 min. HRMS (calc for M+1) 367.0752 (found) 367.0744. IR (KBr, cm<sup>-1</sup>): 3231, 1653, 1341, 1161. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.20-7.23 (m, 1H), 7.47-7.92 (m, 11H), 10.32 (d, 1H, J = 3.6 Hz), 10.90 (d, 1H, J = 3.6 Hz).

## EXAMPLE 13

Quinoline-6-carboxylic acid 2-(phenylsulfonyl)hydrazide




- 36 -

Quinoline-6-carboxylic acid hydrazide (1.0 g, 5.34 mmol) was dissolved in THF (60 mL). N,N-diisopropylethylamine (3.72 mL, 21.36 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.67 mL, 5.34 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 3.5 hours of stirring, the red solution was washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and recrystallized from dichloromethane. The desired product was isolated (0.155 g, 8.9%).

MS: 328.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.42 (1:1 hexane/EtOAc); mp: 211-214°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 1.985, Purity: 90.33%. Analysis (C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S·0.2H<sub>2</sub>O) (calc) C: 58.06, H: 4.08, N: 12.70; (found) C: 57.72, H: 4.04, N: 12.49. IR (KBr, cm<sup>-1</sup>) 3308, 3033, 2817, 1690, 1329, 1164. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.48-7.61 (m, 4H), 7.84 (d, 2H, J = 7.3 Hz), 7.93 (dd, 1H, J = 8.8, 1.7 Hz), 8.01 (d, 1H, J = 8.8 Hz), 8.35 (s, 1H), 8.44 (d, 1H, J = 7.6 Hz), 8.94-8.96 (m, 1H), 10.57 (br, 2H).

## EXAMPLE 14

2-Dibenzofurancarboxylic acid 2-[(2-methylphenyl)sulfonyl]hydrazide



Synthesis of 2-dibenzofurancarboxylic acid 2-[(2-methylphenyl)sulfonyl]hydrazide: Example 14 was synthesized in accordance with the methods of Example 1 except that 3-methylbenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (33% yield).


MS: 381.0 (M+1 for C<sub>20</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>); mp 194-195°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.36 (3:2 hexane/EtOAc); HPLC (C18 column, 70:30 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 98.78%, RT = 4.423 min. HRMS (calc for M+1) 381.0909 (found) 381.0915; IR (KBr, cm<sup>-1</sup>): 3300,

- 37 -

3147, 2964, 1664, 1435, 1157.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  2.29 (s, 3H), 7.35-7.43 (m, 3H), 7.52-7.55 (m, 1H), 7.62-7.74 (m, 4H), 7.84 (d, 1H,  $J = 8.5$  Hz), 8.16 (d, 1H,  $J = 7.8$  Hz), 8.49 (s, 1H), 10.00 (s, 1H), 10.74 (s, 1H).

## EXAMPLE 15

5 2-Dibenzofurancarboxylic acid 2-[(3-methylphenyl)sulfonyl]hydrazide




Synthesis of 2-dibenzofurancarboxylic acid 2-[(3-methylphenyl)sulfonyl]-hydrazide: Example 15 was synthesized in accordance with the methods of Example 1 except that 2-methylbenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (11% yield).

10 MS: 381.0 ( $M+1$  for  $\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_4\text{S}_1$ ); mp 234-235°C. TLC ( $\text{SiO}_2$ )  $R_f = 0.43$  (3:2 hexane/EtOAc); HPLC (C18 column, 70:30  $\text{CH}_3\text{CN}/\text{H}_2\text{O}+0.1\%$  TFA) 97.30%, RT = 4.434 min. HRMS (calc for  $M+1$ ) 381.0909 (found) 381.0911; IR (KBr,  $\text{cm}^{-1}$ ): 3324, 3144, 1657, 1330, 1164.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  2.71 (s, 3H), 7.21-7.55 (m, 5H), 7.68-7.72 (m, 2H), 7.80-7.85 (m, 2H), 8.14 (d, 1H,  $J = 7.8$  Hz), 8.47 (s, 1H), 9.99 (s, 1H), 10.69 (s, 1H).

## EXAMPLE 16

2-Dibenzofurancarboxylic acid 2-[(3-chlorophenyl)sulfonyl]hydrazide



20 Synthesis of 2-dibenzofurancarboxylic acid 2-[(3-chlorophenyl)sulfonyl]-hydrazide: Example 16 was synthesized in accordance with the methods of Example 1


- 38 -

except that 3-chlorobenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (3.0% yield).

MS: 401.0 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Cl<sub>1</sub>); mp 208-209°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.30 (3:2 hexane/EtOAc); HPLC (C18 column, 70:30 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 100%, RT = 4.741 min. HRMS (calc for M+1) 401.0363 (found) 401.0367. IR (KBr, cm<sup>-1</sup>): 3330, 3004, 2804, 1655, 1344, 1192, 1160. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.42 (t, 1H, J = 7.6 Hz), 7.54 (t, 2H, J = 7.8 Hz), 7.68-7.85 (m, 6H), 8.16 (d, 1H, J = 7.6 Hz), 8.50 (s, 1H), 10.31 (s, 1H), 10.82 (s, 1H).

## EXAMPLE 17

2-Dibenzofurancarboxylic acid 2-[(2-chlorophenyl)sulfonyl]hydrazide




Synthesis of 2-dibenzofurancarboxylic acid 2-[(2-chlorophenyl)sulfonyl]hydrazide: Example 17 was synthesized in accordance with the methods of Example 1 except that 2-chlorobenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (25% yield).

MS: 401.1 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Cl<sub>1</sub>); mp 235-236°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.38 (2:1 hexane/EtOAc); HPLC (C18 column, 70:30 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 97.79%, RT = 4.390 min. HRMS (calc for M+1) 401.0363 (found) 401.0359; IR (KBr, cm<sup>-1</sup>): 3338, 3124, 2959, 1719, 1171. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.38-7.73 (m, 7H), 7.85 (d, 1H, J = 8.5 Hz), 7.97 (d, 1H, J = 8.1 Hz), 8.14 (d, 1H, J = 7.6 Hz), 8.51 (s, 1H), 10.15 (s, 1H), 10.75 (s, 1H).

- 39 -

## EXAMPLE 18

Dibenzofuran-2-carboxylic acid 2-(3,5-dichlorophenylsulfonyl)hydrazide



Synthesis of dibenzofuran-2-carboxylic acid 2-(3,5-


5      dichlorophenylsulfonyl)hydrazide: Example 18 was synthesized in accordance with the methods of Example 1 except that 3,5-dichlorobenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (6.7% yield).

MS: 435.0 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Cl<sub>2</sub>); mp: 231-233°C. TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.26(8% MeOH/CH<sub>2</sub>Cl<sub>2</sub>); HPLC (C18 column, 1:1 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 94.51%,

10      RT = 34.421 min. HRMS (calc for M+1) 434.9973 (found) 434.9981. IR (KBr, cm<sup>-1</sup>): 3327, 3079, 2754, 1657, 1349, 1167. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.42-7.46 (m, 2H), 7.52-7.58 (m, 2H), 7.71-7.87 (m, 5H), 8.17-8.22 (m, 1H), 10.51 (br, 1H), 10.87 (br, 1H).

## EXAMPLE 19

15      Dibenzofuran-2-carboxylic acid 2-(3-chloro-2-methyl-phenylsulfonyl)hydrazide



Synthesis of dibenzofuran-2-carboxylic acid 2-(3-chloro-2-methyl-

phenylsulfonyl)hydrazide: Example 19 was synthesized in accordance with the methods of Example 1 except that 3-chloro-2-methyl-benzenesulfonyl chloride was used instead of benzenesulfonyl chloride (4.8% yield).


MS: 415.0 (M+1 for C<sub>20</sub>H<sub>15</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Cl<sub>1</sub>); mp: 192-194°C; TLC (SiO<sub>2</sub>) R<sub>f</sub> = 0.41(8% MeOH/CH<sub>2</sub>Cl<sub>2</sub>); HRMS (calc for M+1): 415.0519 (found): 415.0510. HPLC

- 40 -

(C18 column, 7:3 CH<sub>3</sub>CN/H<sub>2</sub>O+0.1% TFA) 96.38%, RT = 11.222 min. IR (KBr, cm<sup>-1</sup>): 3316, 3042, 1665, 1346, 1191, 1148. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.74 (s, 3H), 7.27 (t, 1H, J = 8.1 Hz), 7.41 (t, 1H, J = 7.6 Hz), 7.53 (t, 1H, J = 7.3 Hz), 7.65-7.73 (m, 3H), 7.80-7.87 (m, 2H), 8.15 (d, 1H, J = 7.3 Hz), 8.48 (s, 1H), 10.27 (br, 1H), 10.73 (br, 1H).

## EXAMPLE 20

Quinoline-6-carboxylic acid 2-(phenylsulfonyl)hydrazide




Quinoline-6-carboxylic acid hydrazide (1.0 g, 5.34 mmol) was dissolved in THF (60 mL). N,N-diisopropylethylamine (3.72 mL, 21.36 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.67 mL, 5.34 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 3.5 hours of stirring, the red solution was washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and recrystallized from dichloromethane. The desired product was isolated (0.155 g, 8.9%).

MS: 328.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.42 (1:1 hexane/EtOAc); mp 211-214°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 1.985, Purity: 90.33%. Analysis (C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S·0.2H<sub>2</sub>O) (calc) C: 58.06, H: 4.08, N: 12.70 (found) C: 57.72, H: 4.04, N: 12.49. IR (KBr, cm<sup>-1</sup>) 3308, 3033, 2817, 1690, 1329, 1164. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.48-7.61 (m, 4H), 7.84 (d, 2H, J = 7.3 Hz), 7.93 (dd, 1H, J = 8.8, 1.7 Hz), 8.01 (d, 1H, J = 8.8 Hz), 8.35 (s, 1H), 8.44 (d, 1H, J = 7.6 Hz), 8.94-8.96 (m, 1H), 10.57 (br, 2H).

- 41 -

## EXAMPLE 21

Quinoline-8-carboxylic acid 2-(phenylsulfonyl)hydrazide




Quinoline-8-carboxylic acid hydrazide (0.3 g, 1.6 mmol) was dissolved in THF (18 mL). N,N-diisopropylethylamine (1.12 mL, 6.41 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.20 mL, 1.6 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 1.5 hours of stirring, the red solution was washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. The desired product was isolated (0.158 g, 30.2%).

MS: 328.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.42 (1:1 hexane/EtOAc); mp: 173-175°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 3.952, Purity: 96.23%. Analysis (C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S) (calc) C: 58.70, H: 4.00, N: 12.84 (found) C: 58.80, H: 4.21, N: 12.66. IR (KBr, cm<sup>-1</sup>) 3190, 1665, 1341, 1172. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.56 (t, 2H, J = 7.8 Hz), 7.63-6.71 (m, 3H), 7.85 (d, 2H, 7.6 Hz), 8.20 (dd, 1H, J = 1.2, 8.1 Hz), 8.32 (d, 1H, J = 7.3 Hz), 8.55 (dd, 1H, J = 1.8, 8.4 Hz), 8.81-8.82 (m, 1H), 10.24 (s, 1H), 12.34 (s, 1H).

## EXAMPLE 22

2-Benzofurancarboxylic acid, 5-chloro-2-[(4-fluorophenyl)]sulfonyl hydrazide



This was run in accordance with the Example 1, Step 3 except that 4-fluorophenyl sulfonyl chloride and 5-chloro-benzofuran-2-carboxylic acid hydrazide were used instead of the benzenesulfonyl chloride and dibenzofuran-2-carboxylic acid hydrazide. The desired product was purified by columning over SiO<sub>2</sub> eluting with

- 42 -

1:2 EtOAc/hexane and triturating the combined and concentrated fractions in diethyl ether (7.9% yield).


MS: 369.0 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl); mp: 229.4-230.5°C. TLC: SiO<sub>2</sub>,

R<sub>f</sub> = 0.75 (EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

5 99.1%, RT = 6.8 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.36 (t, 2H, J = 9.03 Hz), 7.46 (dd, 1H, J = 8.79 and 2.2 Hz), 7.56 (s, 1H), 7.65 (d, 1H, J = 8.79 Hz), 7.85-7.87 (m, 3H), 10.27 (s, 1H), and 11.01 (s, 1H).

## EXAMPLE 23

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-fluorophenyl)sulfonyl]hydrazide



10

This was run in accordance with the Example 22 except that 7-methoxy-benzofuran-2-carboxylic acid hydrazide was used instead of 5-chloro-benzofuran-2-carboxylic acid hydrazide. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined and concentrated

15

MS: 365.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SF); mp: 182.7-183.3°C. TLC: SiO<sub>2</sub>,


R<sub>f</sub> = 0.75 (EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

95.7%, RT = 5.3 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.89 (s, 3H), 7.01 (d, 1H, J = 7.81 Hz), 7.20 (t, 1H, J = 7.87 Hz), 7.25 (d, 1H, J = 7.57 Hz), 7.35 (t, 2H, J = 8.54 Hz), 7.56 (s, 1H), 7.83 (dd, 2H, J = 8.30 and 6.59 Hz), 10.27 (s, 1H), and 11.01 (s, 1H).

20

## EXAMPLE 24

2-Dibenzofurancarboxylic acid 2-[(4-fluorophenyl)sulfonyl]hydrazide



- 43 -

This was run in accordance with Example 1, Step 3 except that 4-fluorophenyl sulfonyl chloride was used instead of benzenesulfonyl chloride (23.1% yield).


MS: 385.0 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>FS); mp: 206.3-207.0°C. TLC: SiO<sub>2</sub>,

R<sub>f</sub> = 0.85 (EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

5 98.0%, RT = 9.72 min. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.08 (t, 2H, J = 8.30 Hz), 7.35 (t, 1H, J = 7.33 Hz), 7.45-7.55 (m, 4H), 7.17 (dd, 1H, J = 8.55 and 1.71 Hz), 7.9-7.96 (m, 3H), 8.19 (d, 1H, J = 5.86 Hz), 8.24 (d, 1H, J = 1.47 Hz).

## EXAMPLE 25

2-Dibenzofurancarboxylic acid 2-[(2-chloro-4-fluorophenyl)sulfonyl]hydrazide



10

This was run in accordance with Example 1, Step 3 except that 2-chloro-4-fluorophenyl sulfonyl chloride was used instead of benzenesulfonyl chloride (4.3% yield).

MS: 419.0 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>FS<sub>2</sub>Cl); mp: 227.5-227.8°C. TLC: SiO<sub>2</sub>,


15 R<sub>f</sub> = 0.85 (EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

97.7%, RT = 13.2 min. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.94 (dd, 1H, J = 10.0 and 2.44 Hz), 7.25 (dd, 1H, J = 7.81 and 1.95 Hz), 7.34 (t, 1H, J = 8.06 Hz), 7.46 (t, 1H, J = 7.33 Hz), 7.55-7.70 (m, 2H), 7.69 (d, 1H, J = 8.55 Hz), 7.89 (d, 1H, J = 7.82 Hz), 7.99-8.08 (m, 3H), 8.23 (s, 1H).

20

## EXAMPLE 26

2-Dibenzofurancarboxylic acid 2-[(3-cyanophenyl)sulfonyl]hydrazide



- 44 -

This was run in accordance with Example 1, Step 3 except that 3-cyanobenzene sulfonyl chloride was used instead of benzenesulfonyl chloride (7.2% yield).


MS: 392.0.0 (M+1 for C<sub>20</sub>H<sub>13</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 234-236°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.20

(EtOAc/hexane; 1:2); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.8%,

RT = 8.12 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.39 (m, 1H), 7.52 (m, 1H), 7.55-7.70 (m, 1H), 7.68-7.72 (m, 3H), 7.81 (dd, 1H, J = 8.79 and 1.95 Hz), 8.06-8.16 (m, 3H), 8.23 (s, 1H), 8.47 (d, 1H, J = 1.46 Hz), 10.41 (s, 1H), 10.84 (s, 1H).

## EXAMPLE 27

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-cyanophenyl)sulfonyl]hydrazide



This was run in accordance with Example 8 except 3-cyanobenzene sulfonyl chloride was used instead of benzenesulfonyl chloride (2.65% yield).


MS: 375.9 (M+1 for C<sub>16</sub>H<sub>10</sub>N<sub>3</sub>O<sub>4</sub>SCl); mp: 208.5-210°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.25

(EtOAc/hexane; 1:2); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 95.3%,

RT = 6.12 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.45 (dd, 1H, J = 8.79 and 1.71 Hz), 7.57 (s, 1H), 7.65 (d, 1H, J = 8.79 Hz), 7.74 (t, 1H, J = 8.06 Hz), 7.84 (s, 1H), 8.10 (t, 2H, J = 8.06 Hz), 8.22 (s, 1H).

## EXAMPLE 28

2-Dibenzofurancarboxylic acid 2-[(2,6-dichlorophenyl)sulfonyl]hydrazide




Dibenzofuran-2-carboxylic acid hydrazide (0.2 g, 0.88 mmol) was dissolved in THF (10 mL). N,N-diisopropylethylamine (0.61 mL, 3.52 mmol) was added, and the reaction was cooled to 0°C. 2,6-Dichloro-benzenesulfonyl chloride (0.22 g, 0.88 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was

- 45 -


allowed to warm to room temperature and stirred overnight. The reaction was then warmed to 40°C and stirred overnight. The solution was diluted with EtOAc (20 mL) and was washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. The desired product was isolated (0.035 g, 9.14%). MS: 437.0 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>Cl<sub>2</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.60 (1:1 hexane/EtOAc); mp: 202-203°C; HRMS: (calc) 434.9973 (found) 434.9977 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>Cl<sub>2</sub>S) HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 17.828, Purity: 98.79%. IR (KBr, cm<sup>-1</sup>) 3494, 3286, 3091, 1673, 1430, 1181, 738. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.39-7.94 (m, 6H), 8.08-8.15 (m, 1H), 8.25 (t, 1H, J = 5.3 Hz), 8.52 (s, 1H), 8.63-8.73 (m, 1H), 10.33 (s, 1H), 10.87 (s, 1H).

## EXAMPLE 29

2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)hydrazide



Step 1. 5-Bromo-benzofuran-2-carboxylic acid hydrazide




This was run in accordance with Example 1, Step 2 except 5-bromo-benzofuran-2-carboxylic acid, which was prepared in accordance with the methods of Perkin, *J. Chem. Soc*, 1951:101-103, was used instead of dibenzofuran-2-carboxylic acid (76% yield).

MS: 254.9 (M+1 for C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Br); mp: 183.5-183.7°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.4%, RT = 1.93 min. Micro: (calc) (C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Br) C: 42.38, H: 2.77, N: 10.98, Br: 31.33; (found) C: 42.33, H: 2.71, N: 10.65, Br: 31.04. <sup>1</sup>H

- 46 -

NMR (DMSO- $d_6$ )  $\delta$  4.57 (s, 2H), 7.45 (d, 1H,  $J = 0.73$  Hz), 7.55 (dd, 1H,  $J = 8.80$  and 1.96 Hz), 7.60 (d, 1H,  $J = 8.80$  Hz), 7.96 (t, 1H,  $J = 1.46$  Hz), and 10.08 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)-hydrazide




This was run in accordance with Example 35 except 5-bromo-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride (28.5% yield).

MS: 396.9 ( $M+1$  for  $C_{15}H_{11}N_2O_4SBr$ ); mp: 168.1-171.1°C. HPLC (C18 column, 1:1 MeCN/ $H_2O$ +0.1% TFA) 96.7%, RT = 6.38 min. Micro: (calc) ( $C_{15}H_{11}N_2O_4SBr$ ) C: 45.59, H: 2.81, N: 7.09, S: 8.11, Br: 20.22; (found) C: 45.76, H: 2.79, N: 7.13, S: 8.26, Br: 20.23.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  7.50-7.56 (m, 3H), 7.58-7.64 (m, 3H), 7.81 (d, 2H,  $J = 7.09$  Hz), 7.81 (d, 1H,  $J = 1.71$  Hz), 10.21 (d, 1H,  $J = 2.8$  Hz), and 11.00 (d, 1H,  $J = 2.8$  Hz).

### EXAMPLE 30

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(2-chloro-4-fluorophenyl)sulfonyl]-hydrazide




This was run in accordance with the Example 8 except that 2-chloro-4-fluorophenyl sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product was purified by column chromatography over  $SiO_2$  eluting with 1:2 EtOAc/hexanes and triturating the combined and concentrated fractions in diethyl ether (2.6% yield).

- 47 -

MS: 404 (M+1 for  $C_{15}H_9N_2O_4SFCl_2$ ); mp: 204.6-206.2°C. TLC:  $SiO_2$ ,  
 $R_f$  = 0.7 (EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/ $H_2O$ +0.1% TFA)  
99.1%, RT = 10.04 min.  $^1H$  NMR ( $DMSO-d_6$ )  $\delta$  7.26-7.31 (m, 1H), 7.43 (dd, 1H,  
J = 9.04 and 2.44 Hz), 7.53 (s, 1H), 7.62 (d, 1H, J = 9.04 Hz), 7.66 (dd, 1H, J = 9.04  
and 2.44 Hz), 7.82 (d, 1H, J = 2.20 Hz), 7.93-7.97 (m, 1H), 10.36 (s, 1H), and 10.97 (s,  
1H).

## EXAMPLE 31

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3,5-dichlorophenyl) sulfonyl]-hydrazide




This was run in accordance with Example 8 except that 3,5-dichlorophenyl  
sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product  
was purified by column chromatography over  $SiO_2$  eluting with 1:2 EtOAc/hexane  
and triturating the combined and concentrated fractions in diethyl ether (4.0% yield).  
MS: 418.9 (M+1 for  $C_{15}H_9N_2O_4SCl_3$ ); mp: 242.5-243.3°C. TLC:  $SiO_2$ ,

$R_f$  = 0.8(EtOAc/hexane; 1:1); HPLC (C18 column, 1:1 MeCN/ $H_2O$ +0.1% TFA)  
99.3%, RT = 17.22 min.  $^1H$  NMR ( $DMSO-d_6$ )  $\delta$  7.47 (dd, 1H, J = 8.79 and 2.20 Hz),  
7.59 (s, 1H), 7.67 (d, 1H, J = 8.79 Hz), 7.75 (d, 2H, J = 1.96 Hz), 7.86 (d, 1H,  
J = 2.20 Hz), 7.97 (s, 1H), 10.63 (s, 1H), and 11.12 (s, 1H).

## EXAMPLE 32

2-Dibenzofurancarboxylic acid, 2-[[3,5-bis(trifluoromethyl)phenyl]sulfonyl]-  
hydrazide



- 48 -

This was run in accordance with Example 1, Step 3 except that 2-(3,5-ditrifluoromethylphenyl)sulfonyl chloride was used instead of benzenesulfonyl chloride (2.3% yield).

MS: 503.0 (M+1 for C<sub>21</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>6</sub>); mp: 244.7-245.5°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.85

(EtOAc/hexane; 1:2); HPLC (cyclohexyl-phenyl column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

98.9%, RT = 13.82 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.41 (t, 1H, J = 7.57 Hz), 7.53 (t, 1H, J = 7.57 Hz), 7.69-7.82 (m, 2H), 8.13 (d, 2H, J = 7.57 Hz), 8.36 (s, 2H), 8.47 (d, 2H, J = 12.21 Hz), 8.65 (s, 1H), 10.72 (s, 1H), 10.99 (s, 1H).

## EXAMPLE 33

2-Benzofurancarboxylic acid, 5-chloro 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide



This was run in accordance with Example 8 except that 3-chloro-4-fluorophenyl sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product was purified by column chromatography over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined and concentrated fractions in diethyl ether (4.2% yield).

MS: 404.9 (M+1 for C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SFCl<sub>2</sub>); mp: 208.2-208.5°C. TLC: SiO<sub>2</sub>,


R<sub>f</sub> = 0.7 (EtOAc/hexane; 1:1); HPLC (cyclohexyl-phenyl column, 1:1

MeCN/H<sub>2</sub>O+0.1% TFA) 99.3%, RT = 10.33 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.45 (dd, 1H, J = 8.79 and 2.20 Hz), 7.55-7.59 (m, 2H), 7.65 (d, 1H, J = 9.03 Hz), 7.66-7.80 (m, 1H), 7.84 (d, 1H, J = 2.20 Hz), 7.96 (dd, 1H, J = 6.84 and 2.20 Hz), 10.46 (s, 1H), and 11.06 (s, 1H).


- 49 -

## EXAMPLE 34

2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide



Step 1. 5-Nitro-benzofuran-2-carboxylic acid hydrazide



5


This was run in accordance with Example 1, Step 2 except 5-nitro-benzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (99% yield).

MS: 222.0 (M+1 for C<sub>9</sub>H<sub>7</sub>N<sub>3</sub>O<sub>4</sub>); HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

97.1%, RT = 1.85 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.60 (s, 2H), 7.67 (s, 1H), 7.84 (d, 1H, J = 9.27 Hz), 8.27 (dd, 1H, J = 2.46 and 9.28 Hz), 8.72 (d, 1H, J = 1.95 Hz), and 10.21 (s, 1H).

10

Step 2. 2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide



This was run in accordance with Example 35 except 5-nitro-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride (31% yield).

15

MS: 362.0 (M+1 for C<sub>15</sub>H<sub>11</sub>N<sub>3</sub>O<sub>6</sub>S); mp: 207.7-208.1°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.6%, RT = 4.38 min. Micro: (calc)

(C<sub>15</sub>H<sub>11</sub>N<sub>3</sub>O<sub>6</sub>S·0.25EtOAc) C: 49.93, H: 3.40, N: 10.92, S: 8.33; (found) C:

20


50.06, H: 3.29, N: 11.29, S: 8.40. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.53 (dt, 2H, J = 8.06 and 1.71 Hz), 7.60-7.65 (m, 1H), 7.79-7.81 (m, 3H), 7.87 (d, 1H, J = 9.28 Hz), 8.30

- 50 -

(dd, 1H, J = 2.44 and 9.28 Hz), 8.76 (d, 1H, J = 2.44 Hz), 10.28 (d, 1H, J = 2.8 Hz), and 11.14 (d, 1H, J = 2.8 Hz).

## EXAMPLE 35

2-Dibenzofurancarboxylic acid 2-[(3-fluorophenyl)sulfonyl]hydrazide



A solution of 3-fluorophenyl sulfonyl chloride (0.19 g, 1 mmol) and dibenzofuran-2-carboxylic acid hydrazide (0.22 g, 1 mmol) was stirred 18 hours at RT in pyridine (10 mL). After the solvent was removed in vacuo, the residue was triturated in water, and dried. Recrystallization of this solid from ethyl acetate provided the title compound, 250 mg, 65% yield.

MS: 385.0 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>FS); mp: 222.6-223.3°C; Analysis:

(C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>FS) (calc) C: 59.37, H: 3.41, N: 7.29, S: 8.34, F: 4.94; (found) C:


59.17, H: 3.45, N: 7.21, S: 8.02, F: 5.03. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1%

TFA) 97.9%, RT = 8.67 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.39 (t, 1H, J = 7.32 Hz),

7.43-7.75 (m, 7H), 7.84 (dd, 1H, J = 8.79 and 1.71 Hz), 8.16 (d, 1H, J = 7.82 Hz), 8.50 (d, 1H, J = 1.22 Hz), 10.28 (d, 1H, J = 2.8 Hz), 10.812 (d, 1H, J = 2.8 Hz).

## EXAMPLE 36

2-Dibenzofurancarboxylic acid 2-[(5-fluoro-2-methylphenyl)sulfonyl]hydrazide



This was run in accordance with Example 35 except 5-fluoro-2-methylphenylsulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride (55.3% yield).

MS: 399.0 (M+1 for C<sub>20</sub>H<sub>15</sub>N<sub>2</sub>O<sub>4</sub>FS); mp: 210.1-210.5°C; Analysis:


(C<sub>20</sub>H<sub>15</sub>N<sub>2</sub>O<sub>4</sub>FS) (calc) C: 59.62, H: 3.88, N: 6.95, S: 7.96, F: 4.471; (found) C:

- 51 -

59.54, H: 3.74, N: 6.99, S :7.78, F: 4.84. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.2%, RT = 9.25 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.65 (s, 3H), 7.32-7.39 (m, 1H), 7.45 (dd, 2H, J = 7.57 and 1.22 Hz), 7.55 (dd, 1H, J = 7.57 and 1.22 Hz), 7.60 (dd, 1H, J = 9.04 and 2.69 Hz), 7.72 (t, 1H, J = 8.79 Hz), 7.83 (dd, 1H, J = 8.54 and 1.46 Hz), 8.15 (d, 1H, J = 7.58 Hz), 8.50 (s, 1H), 10.24 (d, 1H, J = 2.8 Hz), 10.76 (d, 1H, J = 2.8 Hz).

## EXAMPLE 37

2-Dibenzofurancarboxylic acid 2-[(3,4-difluorophenyl) sulfonyl]hydrazide




This was run in accordance with Example 35 except 3,4-difluorophenylsulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride (42.4% yield), and the material was purified through column chromatography eluting with 1:2 ethyl acetate/hexanes.

MS: 402.9 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>F<sub>2</sub>S); mp: 210.1-210.5°C; Analysis:

(C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>F<sub>2</sub>S) (calc) C: 56.72, H: 3.01, N: 6.96, S: 7.97, F: 9.44; (found) C: 56.64, H: 3.00, N: 6.88, S: 7.71, F: 9.48. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.2%, RT = 11.06 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.37-7.41 (m, 1H), 7.52 (dd, 2H, J = 7.33 and 1.22 Hz), 7.55-7.73 (m, 3H), 7.82-7.90 (m, 2H), 8.14 (d, 1H, J = 7.82 Hz), 8.15 (d, 1H, J = 7.58 Hz), 8.49 (d, 1H, J = 1.22 Hz), 10.3 (s, 1H), 10.8 (s, 1H).

## EXAMPLE 38

2-Dibenzofurancarboxylic acid 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide




- 52 -

This was run in accordance with Example 35 except 3-chloro-4-fluorophenylsulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride (28.5% yield), and the material was purified through column chromatography eluting with 1:2 ethyl acetate/hexanes.

5 MS: 418.9 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>ClFS); mp: 210.8-211.6°C; Analysis: (C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>ClFS) (calc) C: 54.49, H: 2.89, N: 6.69, S: 7.66, F: 4.54; (found) C: 53.97, H: 3.02, N: 6.53, S: 7.41, F: 4.64. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.9%, RT = 15.05 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.40 (dd, 1H, J = 7.57 and 0.73 Hz), 7.49-7.57 (m, 2H), 7.68 (d, 1H, J = 8.30 Hz), 7.71 (d, 1H, J = 8.30 Hz),  
10 7.79-7.84 (m, 2H), 7.99 (dd, 1H, J = 6.84 and 1.95 Hz), 8.14 (d, 1H, J = 7.57 Hz), 8.48 (d, 1H, J = 0.68 Hz), 10.33 (s, 1H), 10.81 (s, 1H).

## EXAMPLE 39

2-Dibenzofurancarboxylic acid 2-[(3-methoxyphenyl)sulfonyl]hydrazide




15 This was run in accordance with Example 35 except 3-methoxyphenylsulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride (63.1% yield).

MS: 397.0 (M+1 for C<sub>20</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 199.5-200.5°C; Analysis: (C<sub>20</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S) (calc) C: 60.60, H: 4.07, N: 7.07, S: 8.09; (found) C: 60.24, H: 3.96, N: 6.95, S: 7.86. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%,  
20 RT = 8.60 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.72 (s, 3H), 7.14-7.17 (m, 1H), 7.34-7.39 (m, 1H), 7.40-7.43 (m, 3H), 7.55 (t, 1H, J = 5.86 Hz), 7.84 (dd, 1H, J = 9.08 and 2.0 Hz), 8.15 (d, 1H, J = 6.84 Hz), 8.50 (s, 1H), 10.07 (d, 1H, J = 2.80 Hz), and 10.75 (d, 1H, J = 2.80 Hz).

- 53 -

## EXAMPLE 40

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-fluorophenyl)sulfonyl] hydrazide




This was run in accordance with the Example 35 except that 5-chloro-  
5 benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-  
2-carboxylic acid hydrazide. The desired product was purified by columning over  
SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined and concentrated  
fractions in diethyl ether (46.2% yield).

MS: 368.9 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SFCl); mp: 174.4-174.7°C; Micro: (calc)  
10 (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SFCl) C: 48.86, H: 2.73, N: 7.60, S: 8.69, F: 5.15; (found) C: 48.56, H:  
2.61, N: 7.45, S: 8.64, F: 5.19. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)  
98.9%, RT = 7.73 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.46 (dd, 2H, J = 9.04 and 2.2 Hz),  
7.50-7.67 (m, 5H), 7.85 (d, 1H, J = 1.96 Hz), 10.41 (s, 1H), and 11.05 (s, 1H).

## EXAMPLE 41

2-Benzofurancarboxylic acid, 5-chloro-2-[(3,4-difluorophenyl)sulfonyl]hydrazide



This was run in accordance with Example 40 except that 3,4-difluorophenyl  
sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired  
product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (51.7%  
20 yield).


MS: 386.9 (M+1 for C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>2</sub>Cl); mp: 193.8-195°C; Micro: (calc)  
(C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>2</sub>Cl) C: 46.58, H: 2.35, N: 7.25, S: 8.29, F: 9.82; (found) C: 46.58, H:  
2.44, N: 7.08, S: 8.18, F: 9.54. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)  
97.5%, RT = 8.34 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.46 (dd, 1H, J = 9.04 and 2.2 Hz),

- 54 -

7.57 (s, 1H), 7.61 (dd, 1H, J = 10.25 and 2.68 Hz), 7.65-7.83 (m, 2H), 7.84-7.88 (m, 3H), 10.45 (s, 1H), and 11.06 (s, 1H).

## EXAMPLE 42

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-chlorophenyl)sulfonyl]hydrazide




This was run in accordance with Example 40 except that 2-chlorophenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (51.9% yield).

MS: 384.9 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 200.3-200.5°C; Micro: (calc) (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 46.77, H: 2.62, N: 7.27, S: 8.32, Cl: 18.41; (found) C: 47.09, H: 2.98, N: 6.58, S: 7.57, Cl: 18.62. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.2%, RT = 7.49 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.41 (dd, 1H, J = 6.59 and 1.71 Hz), 7.45 (dd, 1H, J = 8.79 and 2.2 Hz), 7.54 (s, 1H), 7.58-7.65 (m, 3H), 7.82 (d, 1H, J = 1.96), 7.92 (dd, 1H, J = 9.03 and 1.22 Hz), 10.30 (s, 1H), and 10.98 (s, 1H).

## EXAMPLE 43

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chlorophenyl) sulfonyl]hydrazide




This was run in accordance with Example 40 except that 3-chlorophenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (42.4% yield).

- 55 -

MS: 384.9 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 202.9-205.2°C; Micro: (calc) for (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 46.77, H: 2.62, N: 7.27, S: 8.32, Cl: 18.41; (found): C: 46.71, H: 2.43, N: 7.18, S: 8.25, Cl: 18.35; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%, RT = 9.3 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.46 (dd, 1H, J = 8.79 and 2.19 Hz), 7.55 (d+s, 2H, J = 7.82 Hz), 7.54 (s, 1H), 7.70-7.75 (m, 2H), 7.80 (t, 1H, J = 1.96), 7.85 (d, 1H, J = 2.20 Hz), 10.44 (s, 1H), and 11.07 (s, 1H).

## EXAMPLE 44

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-bromophenyl)sulfonyl]hydrazide




This was run in accordance with Example 40 except that 2-bromophenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (59.8% yield).

MS: 426.9 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SBrCl); mp: 206.4-207.3°C; Micro: (calc) (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SBrCl·0.5H<sub>2</sub>O) C: 41.68, H: 2.62, N: 5.96; (found) C: 41.07, H: 2.53, N: 6.39. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 93.7%, RT = 8.17 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.44-7.52 (m, 3H), 7.55 (d, 1H, J = 0.73 Hz), 7.64 (d, 1H, J = 8.80 Hz), 7.79-7.82 (m, 1H), 7.84 (d, 1H, J = 1.95 Hz), 7.97-7.99 (m, 1H), 10.30 (br s, 1H), and 10.95 (br s, 1H).

## EXAMPLE 45

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-bromophenyl)sulfonyl]hydrazide



- 56 -

This was run in accordance with Example 40 except that 3-bromophenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by recrystallization in ethyl acetate (58.3% yield).

MS: 428.9 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SBrCl); mp: 218.1-220.0°C; Micro: (calc) for

(C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SBrCl) C: 42.06, H: 2.36, N: 6.54, S: 7.47; (found) C: 41.99, H:

2.31, N:6.51, S: 7.21. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 95.8%,

RT = 10.17 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.45 (t, 2H, J = 8.06 Hz), 7.57 (s, 1H), 7.66


(d, 1H, J = 9.04 Hz), 7.76 (dd, 1H, J = 1.71 and 0.98 Hz), 7.78 (dd, 1H, J = 1.71 and

0.98 Hz), 7.85 (s, 1H), 7.92 (t, 1H, J = 1.71 Hz), 10.44 (d, 1H, J = 2.80 Hz), and

11.08 (d, 1H, J = 2.80 Hz).

## EXAMPLE 46

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-methylphenyl)sulfonyl]hydrazide



This was run in accordance with Example 40 except that 2-methylphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (63.0% yield).

MS: 364.9 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl); mp: 190.5-192.0°C; Micro: (calc)

(C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl·0.5 EtOAc) C: 52.88, H: 4.19, N: 6.85, S:7.84; (found): C:

52.90, H: 3.91, N: 7.00, S: 8.03. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

98.8%, RT = 13.1 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.66 (s, 3H), 7.25 (t, 1H, J = 7.57 Hz),


7.35 (d, 1H, J = 7.57 Hz), 7.43-7.49 (m, 2H), 7.52 (s, 1H), 7.63 (d, 1H, J = 9.04 Hz),

7.78 (d, 1H, J = 6.84 Hz), 7.83 (d, 1H, J = 2.20 Hz), 10.12 (s, 1H), and 10.91 (s, 1H).

- 57 -

## EXAMPLE 47

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methylphenyl)sulfonyl]hydrazide



This was run in accordance with Example 40 except that 3-methylphenyl sulfonyl chloride was used instead of 4-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (83.0% yield).

MS: 364.9 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl); mp: 201.2-201.6°C; Micro: (calc)


(C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl) C: 52.68, H: 3.59, N: 7.68, S: 8.74. (found) C: 52.61, H: 3.61, N:

7.57, S: 8.67. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.8%, RT = 12.3

min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.31 (s, 3H), 7.37-7.48 (m, 3H), 7.56 (d, 1H, J = 0.73 Hz), 7.59 (d, 1H, J = 7.33 Hz), 7.65 (t, 2H, J = 11.23 Hz), 7.85 (d, 1H, J = 2.20 Hz), 10.17 (s, 1H), and 10.97 (s, 1H).

## EXAMPLE 48

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methoxyphenyl)sulfonyl] hydrazide



This was run in accordance with Example 40 except that 3-methoxyphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes (68.2% yield).

MS: 380.9 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SCl); mp: 204.3-204.6°C; Micro: (calc)

(C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SCl-0.3H<sub>2</sub>O) C: 49.76, H: 3.43, N: 7.13, S: 8.19, Cl: 9.17; (found) C:

49.75, H: 3.55, N: 7.25, S: 8.30, Cl: 9.18; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1%


TFA) 98.5%, RT = 6.35 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.74 (s, 3H), 7.18 (dd, 1H,

- 58 -

J = 7.08 and 0.98 Hz), 7.30 (s, 1H), 7.36 (t, 1H, J = 7.08 Hz), 7.41 (d, 1H, J = 7.82 Hz), 7.47 (d, 1H, J = 2.20 Hz), 7.55 (d, 1H, J = 0.74 Hz), 7.65 (d, 1H, J = 8.79 Hz), 7.85 (d, 1H, J = 1.95 Hz), 10.17 (s, 1H), and 10.71 (s, 1H).

## EXAMPLE 49

5 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chloro-2-methylphenyl)sulfonyl]-hydrazide




This was run in accordance with Example 40 except that 3-chloro-2-methylphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexanes and triturating the combined pure fractions in ethyl ether (62.7% yield).

MS: 398.9 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 198.2-198.5°C; Micro: (calc) (C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 48.13, H: 3.03, N: 7.02, S: 8.03, Cl: 17.76; (found) C: 48.09, H: 3.11, N: 6.90, S: 8.02, Cl: 17.80; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 93.25%, RT = 7.65 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.70 (s, 3H), 7.28 (t, 1H, J = 7.81 Hz), 7.45 (dd, 1H, J = 8.79 and 2.20 Hz), 7.54 (s, 1H), 7.64 (d, 1H, J = 9.04 Hz), 7.69 (d, 1H, J = 7.81 Hz), 7.80 (d, 1H, J = 7.81 Hz), 7.83 (d, 1H, J = 2.20 Hz), 10.39 (s, 1H), and 10.96 (s, 1H).

20

## EXAMPLE 50

2-Benzofurancarboxylic acid, 5-chloro-2-[(5-fluoro-2-methylphenyl)sulfonyl]-hydrazide




- 59 -

This was run in accordance with Example 40 except that 5-fluoro-2-methylphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined pure fractions in ethyl ether (52.2% yield).

MS: 382.9 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SFCl); mp: 209.6-211.2°C; Micro: (calc) (C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SFCl) C: 50.20, H: 3.16, N: 7.32, S: 8.38, F: 4.96; (found) C: 49.83, H: 3.04, N: 7.17, S: 8.15, F: 4.93; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.8%, RT = 11.77 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.61 (s, 3H), 7.34-7.43 (m, 2H), 7.45 (dd, 1H, J = 8.79 and 2.20 Hz), 7.55 (s, 1H), 7.57 (d, 1H, J = 2.68 Hz), 7.65 (d, 1H, J = 8.79 Hz), 7.84 (d, 1H, J = 2.2 Hz), 10.39 (s, 1H), and 10.96 (s, 1H).

## EXAMPLE 51

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-(trifluoromethyl)phenyl)sulfonyl]-hydrazide




This was run in accordance with Example 40 except that 2-trifluoromethylphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane (60.3% yield).

MS: 419.9 (M+1 for C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>3</sub>Cl); mp: 192.0-193.4°C; Micro: (calc) (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>3</sub>Cl) C: 45.89, H: 2.41, N: 6.69, S: 7.66, F: 13.61; (found) C: 46.16, H: 2.41, N: 6.60, S: 7.38, F: 13.30; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.2%, RT = 9.63 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.44 (dd, 1H, J = 9.04 and 2.20 Hz), 7.53 (s, 1H), 7.63 (d, 1H, J = 8.79 Hz), 7.75-7.82 (m, 2H), 7.84 (d, 1H, J = 1.95 Hz), 7.93 (dd, 1H, J = 2.2 and 9.03 Hz), 8.13 (dd, 1H, J = 2.44 and 6.59 Hz), 10.19 (s, 1H), and 11.06 (s, 1H).

- 60 -

## EXAMPLE 52

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethoxy)phenyl]sulfonyl]-hydrazide




This was run in accordance with Example 40 except that 2-trifluoromethoxyphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by recrystallizing in ethyl acetate and then columning the resulting solid over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane (34.6% yield).

MS: 434.9 (M+1 for C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>Cl); mp: 218.2-219.8°C; Micro: (calc) (C<sub>16</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>Cl) C: 44.20, H: 2.32, N: 6.44, S: 7.37, F: 13.11; (found) C: 44.10, H: 2.46, N: 6.27, S: 7.23, F: 13.22; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.4%, RT = 11.9 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.54-7.58 (m, 2H), 7.62 (d, 1H, J = 8.31 Hz), 7.65 (d, 1H, J = 0.97 Hz), 7.76 (d, 1H, J = 9.03 Hz), 7.84 (dt, 1H, J = 1.71 and 8.31 Hz), 7.96 (d, 1H, J = 1.96 Hz), 8.02 (dd, 1H, J = 1.70 and 8.06 Hz), 10.46 (s, 1H), and 11.15 (s, 1H).

## EXAMPLE 53

2-Dibenzofurancarboxylic acid, 2-[(4-N,N-dimethylaminophenyl)-sulfonyl]hydrazide hydrochloride salt



Synthesis of 2-Dibenzofurancarboxylic acid, 2-[(4-N,N-dimethylaminophenyl)sulfonyl] hydrazide hydrochloride salt: Example 53 was synthesized in accordance with the methods of Example 167, Step 2, except that 2-dibenzofurancarboxylic acid, 2-[(4-N,N-dimethylaminophenyl)sulfonyl]-hydrazide


- 61 -

was used instead of 2-dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl)sulfonyl]hydrazide (yield 98%).

MS: 409.1 (M for C<sub>21</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 187.6-188.8°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 88.97%, RT = 8.3 min.; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ 3.15 (s, 6H), 7.00 (t, 1H, J = 7.1 Hz), 7.18 (d, 1H, J = 8.1 Hz), 7.40-7.80 (m, 6H), 7.92 (d, 1H, J = 7.8 Hz), 8.18 (d, 1H, J = 7.4 Hz), 8.55 (s, 1H), 10.75 (s, 2H).

## EXAMPLE 54

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide




This was run in accordance with Example 40 except that 7-methoxy-benzofuran-2-carboxylic acid hydrazide was used instead of 5-chlorobenzofuran-2-carboxylic acid hydrazide, and 2-trifluoromethylphenyl sulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined and concentrated fractions in diethyl ether (41.6% yield).

MS: 415.1 (M+1 for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>); mp: 156.8-157.6°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 6.49 min. Micro: (calc) (C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>) C: 49.28, H: 3.16, N: 6.76, S: 7.74, F: 13.75; (found) C: 49.31, H: 3.12, N: 6.67, S: 7.70, F: 13.76; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.89 (s, 3H), 7.02 (d, 1H, J = 7.81 Hz), 7.21 (t, 1H, J = 7.58 Hz), 7.26 (d, 1H, J = 7.08 Hz), 7.54 (s, 1H), 7.76-7.83 (m, 3H), 7.93 (d, 1H, J = 7.08 Hz), 8.14 (d, 1H, J = 6.84 Hz), 10.12 (s, 1H), and 10.89 (s, 1H).

- 62 -

## EXAMPLE 55


2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methylphenyl) sulfonyl] hydrazide




This was run in accordance with Example 54 except 3-methylphenyl sulfonyl chloride was used instead of 2-trifluoromethylphenyl sulfonyl chloride. The desired product was purified by columning over SiO<sub>2</sub> eluting with 1:2 EtOAc/hexane and triturating the combined and concentrated fractions in diethyl ether (33.3% yield). MS: 361.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 170.8-171.8°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%, RT = 5.08 min. Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S) C: 56.66 H: 4.48, N: 7.77, S: 8.90; (found) C: 56.45, H: 4.49, N: 7.55, S: 8.68; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.31 (s, 3H), 3.90 (s, 3H) 7.02 (d, 1H, J = 7.57 Hz), 7.20 (t, 1H, J = 7.81 Hz), 7.26 (d, 1H, J = 7.82 Hz), 7.36-7.43 (m, 2H), 7.55 (s, 1H), 7.57-7.62 (m, 2H), 10.12 (s, 1H), and 10.89 (s, 1H).

## EXAMPLE 56

2-Benzofurancarboxylic acid, 6-methoxy-, 2-(phenylsulfonyl)hydrazide



Step 1. 6-Methoxy-benzofuran-2-carboxylic acid hydrazide




The hydrazide was prepared in accordance with Example 1, Step 2 except 6-methoxy-benzofuran-2-carboxylic acid, prepared in accordance with the methods of Tanaka et al., *J. Amer. Chem. Soc.*, 1951:872, was used instead of the dibenzofuran-2-carboxylic acid (yield 58.9%).

- 63 -

MS: 207.0 (M+1 for C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>); mp: 124.1-125.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%, RT = 1.67 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.79 (s, 3H), 4.49 (s, 2H), 6.90 (dd, 1H, J = 8.54 and 2.19 Hz), 7.16 (d, 1H, J = 1.95 Hz), 7.40 (d, 1H, J = 0.73 Hz), 7.58 (d, 1H, J = 8.79 Hz), and 9.84 (s, 1H).

5 Step 2. 2-Benzofurancarboxylic acid, 6-methoxy-, 2-(phenylsulfonyl) hydrazide




This was run in accordance with Example 35 except 6-methoxy-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and benzenesulfonyl chloride was used instead of 3-fluorophenyl sulfonyl chloride. The desired product was purified by recrystallization in ethyl acetate/methanol (66.5% yield).

10 MS: 347.0 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 224.9-225.8°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 4.18 min. Micro: (calc) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>S) C: 55.48, H: 4.07, N: 8.09; (found) C: 55.37, H: 4.00, N: 8.07. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.78 (s, 3H) 6.92 (dd, 1H, J = 8.55 and 2.20 Hz), 7.16 (d, 1H, J = 1.95 Hz), 7.50-7.53 (m, 3H), 7.59-7.61 (m, 2H), 7.57-7.62 (m, 2H), 7.79 (dd, 1H, J = 7.08 and 1.46 Hz), 10.12 (s, 1H), and 10.77 (s, 1H).


#### EXAMPLE 57

20 2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl)-hydrazide



- 64 -

Step 1. 5-Methoxy-3-(1-methylethoxy)-benzofuran-2-carboxylic acid hydrazide




The hydrazide was prepared in accordance with Example 1, Step 2 except that 5-methoxy-3-(1-methylethoxy)-benzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (yield 59.9%).

MS: 265.3 (M+1 for C<sub>13</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>); HPLC (C18 column, 1:1

MeCN/H<sub>2</sub>O+0.1% TFA) 95.5%, RT = 1.88 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.39 (d, 6H, J = 6.10 Hz), 3.88 (s, 3H), 4.58 (d, 2H, J = 3.18 Hz), 5.00 (quintet, 1H, J = 6.11 Hz), 7.12 (dd, 1H, J = 9.09 and 2.49 Hz), 7.19 (d, 1H, J = 2.45 Hz), 7.53 (d, 1H, J = 9.03 Hz), and 9.29 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl) hydrazide



This was run in accordance with Example 35 except 3-isopropoxy-5-methoxy-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by column chromatography with ethyl acetate/hexanes (45.6% yield).


MS: 405.3 (M+1 for C<sub>19</sub>H<sub>20</sub>N<sub>2</sub>O<sub>6</sub>S); mp: 127-129°C. HPLC (C18 column, 1:1

MeCN/H<sub>2</sub>O+0.1% TFA) 98.6%, RT = 6.14 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.29 (s, 6H), 3.76 (s, 3H), 4.64 (quintet, 1H), 7.03-7.06 (s+m, 2H), 7.43 (d, 1H, J = 8.54 Hz), 7.52 (t, 2H, J = 8.06 Hz), 7.59-7.63 (m, 1H), 7.80 (dd, 2H, J = 7.32 and 1.47 Hz), 10.00 (s, 1H), and 10.06 (s, 1H).


- 65 -

## EXAMPLE 58

2-Benzofurancarboxylic acid, 3, 5-dimethoxy-, 2-(phenylsulfonyl)hydrazide



Step 1. 3,5-Dimethoxy-benzofuran-2-carboxylic acid hydrazide




5

The hydrazide was prepared in accordance with Example 1, Step 2 except that 3,5-dimethoxy-benzofuran-2-carboxylic acid, which was prepared in accordance with the methods of Lamotte et al., *Eur. J. Med Chem. Chim. Ther.*, 1986:379-383, was used instead of dibenzofuran-2-carboxylic acid (yield 89.9%).

10

MS: 237.0 (M+1 for  $C_{11}H_{12}N_2O_4$ ); mp: 176.2-179.1°C.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  3.78 (s, 3H), 4.11 (s, 3H), 4.83 (s, 2H), 7.02 (dd, 1H, J = 9.04 and 2.69 Hz), 7.21 (d, 1H, J = 2.45 Hz), 7.43 (d, 1H, J = 9.03 Hz), and 9.25 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 3,5-dimethoxy-, 2-(phenylsulfonyl)-hydrazide



15

This was run in accordance with Example 56 except 3,5-dimethoxy-benzofuran-2-carboxylic acid hydrazide was used instead of 6-methoxy-benzofuran-2-carboxylic acid hydrazide.

MS: 377.0 (M+1 for  $C_{17}H_{16}N_2O_6S$ ); mp: 172.2-173.8°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.3%, RT = 4.18 min. Micro: (calc)

20


( $C_{17}H_{16}N_2O_6S \cdot 0.5H_2O$ ) C: 52.77 H: 4.43, N: 7.24; (found) C: 53.03, H: 4.22, N: 7.09.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  3.78 (s, 3H), 3.87 (s, 3H), 7.06 (dd, 2H, J = 9.04 and 2.69 Hz), 7.18 (d, 1H, J = 2.45 Hz), 7.44 (d, 1H, J = 9.04 Hz), 7.52 (t, 2H, J = 7.09

- 66 -

Hz), 7.61 (t, 1H, J = 7.33 Hz), 7.80 (d, 2H, J = 7.33 Hz), 10.04 (d, 1H, J = 3.20 Hz), and 10.18 (d, 1H, J = 3.20 Hz).


## EXAMPLE 59

2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)hydrazide



5

Step 1. 3,6-Dimethoxy-benzofuran-2-carboxylic acid hydrazide




The hydrazide was prepared in accordance with Example 1, Step 2 except that 3,6-dimethoxy-benzofuran-2-carboxylic acid, which was prepared in accordance with Lele, *J. Sci. Ind. Res.*, 1955:101-103, was used instead of dibenzofuran-2-carboxylic acid (yield 85.9%).

10

MS: 237.0 (M+1 for C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>); mp: 178.9-181.5°C. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.79 (s, 3H), 4.13 (s, 3H), 4.44 (s, 2H), 6.90 (dd, 1H, J = 8.80 and 2.20 Hz), 7.09 (d, 1H, J = 1.95 Hz), 7.71 (d, 1H, J = 8.79 Hz), and 8.97 (s, 1H).

15

Step 2. 2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide



This was run in accordance with Example 56 except 3,6-dimethoxy-benzofuran-2-carboxylic acid hydrazide was used instead of 6-methoxy-benzofuran-2-carboxylic acid hydrazide. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (37.4% yield).

20


MS: 377.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S); mp: 174.8-175.2°C; HPLC (C<sub>18</sub> column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.7%, RT = 4.20 min. Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S·0.2H<sub>2</sub>O) C: 53.72, H: 4.35, N: 7.37; (found) C: 53.77, H: 4.02, N:

- 67 -


7.11.  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$  3.79, (s, 3H), 4.02 (s, 3H), 6.90 (dd, 2H,  $J = 8.80$  and 2.20 Hz), 7.08 (d, 1H,  $J = 1.95$  Hz), 7.52 (t, 2H,  $J = 7.33$  Hz), 7.61 (t, 1H,  $J = 7.33$  Hz), 7.72 (d, 1H,  $J = 8.79$  Hz), 7.81 (d, 2H,  $J = 7.33$  Hz), 9.85 (s, 1H), and 9.99 (s, 1H).

## EXAMPLE 60

2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)hydrazide



Step 1. 4,6-Dimethoxy-benzofuran-2-carboxylic acid hydrazide




The hydrazide was prepared in accordance with Example 1, Step 2 except that 4,6-dimethoxy-benzofuran-2-carboxylic acid, which was prepared in accordance with Reichstein et al., *Helv. Chim. Acta*, 1935:816, 828, was used instead of dibenzofuran-2-carboxylic acid (yield 50%).

MS: 237.0 ( $M+1$  for  $\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_4$ ); mp: 185.8-186.6°C. HPLC (C18 column, 1:1

$\text{MeCN}/\text{H}_2\text{O}+0.1\%$  TFA) 99.7%, RT = 1.72 min.  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$  3.78 (s, 3H), 3.86 (s, 3H), 4.51 (s, 2H), 6.42 (d, 1H,  $J = 1.96$  Hz), 7.37 (t, 1H,  $J = 0.73$  Hz), 7.37 (d, 1H,  $J = 0.73$  Hz), and 9.78 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)hydrazide



This was run in accordance with Example 56 except 4,6-dimethoxy-benzofuran-2-carboxylic acid hydrazide was used instead of 6-methoxy-benzofuran-

- 68 -


2-carboxylic acid hydrazide. The desired product was purified by recrystallization in ethyl acetate/methanol (83.8% yield).

MS: 377.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S); mp: 230.6-231.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.1%, RT = 5.73 min. Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S) C:


54.25, H: 4.28, N: 7.44, S: 8.52; (found) C: 54.34, H: 4.06, N: 7.38, S: 8.60. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.77 (s, 3H), 3.84 (s, 3H), 6.42 (d, 1H, J = 1.95 Hz), 6.76 (d, 1H, J = 0.73 Hz), 7.50 (dd, 2H, J = 8.06 and 1.95 Hz), 7.61 (t, 1H, J = 7.33 Hz), 7.78 (dd, 2H, J = 7.08 and 1.46 Hz), 10.67 (s, 1H), and 10.67 (s, 1H).

## EXAMPLE 61

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl)hydrazide



Step 1. 3,7-Dimethoxy-benzofuran-2-carboxylic acid hydrazide




The hydrazide was prepared in accordance with Example 1, Step 2 except that 3,7-dimethoxy-2-benzofurancarboxylic acid, prepared in accordance with the methods of Lamotte et al., *Eur. J. Med Chem. Chim. Ther.*, 1986:379-383, was used instead of dibenzofuran-2-carboxylic acid (yield 71%).

MS: 237.0 (M+1 for C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>); mp: 172.1-173.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.7%, RT = 1.64 min. Micro: (calc)

(C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>·0.2H<sub>2</sub>O) C: 55.09, H: 5.21, N: 11.68; (found) C: 55.08, H: 5.04, N: 11.16. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.91(s, 3H), 4.11 (s, 3H), 4.49 (s, 2H), 7.04 (d, 1H, J = 7.81 Hz), 7.20 (t, 1H, J = 7.81 Hz), 7.36 (d, 1H, J = 7.81 Hz), and 9.15 (s, 1H).

- 69 -

Step 2. 2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl-hydrazide




This was run in accordance with Example 56 except 3,7-dimethoxy-benzofuran-2-carboxylic acid hydrazide was used instead of except 6-methoxy-benzofuran-2-carboxylic acid hydrazide. The desired product was purified by

recrystallization in ethyl acetate/hexanes (66.5% yield).  
 MS: 377.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S); mp: 177.9-179.6°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.1%, RT = 4.35 min. Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S) C: 54.25, H: 4.28, N: 7.44, S: 8.52; (found) C: 54.01, H: 4.23, N: 6.98, S: 8.64. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.91 (s, 3H), 4.00 (s, 3H), 7.07 (d, 1H, J = 7.82 Hz), 7.20 (t, 1H, J = 8.06 Hz), 7.35 (d, 1H, J = 7.82 Hz), 7.62 (t, 2H, J = 8.06 Hz), 7.59-7.62 (m, 1H), 7.81 (d, 2H, J = 7.33 Hz), 10.05 (d, 1H, J = 3.2 Hz), and 10.13 (d, 1H, J = 3.2 Hz).

#### EXAMPLE 62

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(2-trifluoromethylphenyl)-sulfonyl]hydrazide



This was run in accordance with Example 61 except 2-trifluoromethylphenyl sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (46.5% yield).


MS: 445.0 (M+1 for C<sub>18</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SF<sub>3</sub>); mp: 190.6-192.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.1%, RT = 6.17 min. Micro: (calc) (C<sub>18</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SF<sub>3</sub>) C: 48.65, H: 3.40, N: 6.30, S: 7.22; (found) C: 48.59, H: 3.34, N: 6.17, S: 7.26. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.90 (s, 3H), 3.98 (s, 3H), 7.07 (d, 1H, J = 7.33 Hz), 7.21 (t, 1H,

- 70 -

J = 8.06 Hz), 7.35 (d, 1H, J = 7.08 Hz), 7.70-7.81 (m, 2H), 7.94 (t, 2H, J = 4.15 Hz), 8.16 (t, 2H, J = 4.88 Hz), 10.04 (d, 1H, J = 2.0 Hz), and 10.31 (d, 1H, J = 2.0 Hz).

## EXAMPLE 63


2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(4-fluorophenyl)sulfonyl]-hydrazide



This was run in accordance with Example 61 except 4-fluorophenyl sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (72% yield). MS: 395.0 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SF); mp: 190.9-191.1°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 96.8%, RT = 4.90 min. Micro: (calc) (C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SF) C: 51.77, H: 3.83, N: 7.10, S: 8.13, F: 4.82; (found) C: 51.73, H: 3.73, N: 7.02, S: 8.04, F: 4.88. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.91 (s, 3H), 4.01 (s, 3H), 7.07 (d, 1H, J = 8.06 Hz), 7.21 (t, 1H, J = 8.06 Hz), 7.37 (t, 3H, J = 9.04 Hz), 7.84-7.88 (m, 2H), 10.12 (s, 1H), and 10.20 (s, 1H).

## EXAMPLE 64

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(3-methylphenyl)sulfonyl]-hydrazide




This was run in accordance with Example 61 except 3-methylphenyl sulfonyl chloride was used instead of benzenesulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (70% yield). MS: 391.0 (M+1 for C<sub>18</sub>H<sub>18</sub>N<sub>2</sub>O<sub>6</sub>S); mp: 179.4-180.9°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.8%, RT = 5.22 min. Micro: (calc) (C<sub>18</sub>H<sub>18</sub>N<sub>2</sub>O<sub>6</sub>S·0.7H<sub>2</sub>O) C: 53.55, H: 4.84, N: 6.94, S: 7.94; (found) C: 53.61, H:

- 71 -


4.37, N: 6.88, S: 8.19.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  3.90 (s, 3H), 4.01 (s, 3H), 7.07 (d, 1H,  $J = 8.06$  Hz), 7.21 (t, 1H,  $J = 8.06$  Hz), 7.36 (d, 1H,  $J = 8.06$  Hz), 7.38-7.40 (m, 1H), 7.42 (s, 1H), 7.60 (d, 1H,  $J = 6.83$  Hz), 7.64 (s, 1H), 10.0 (s, 1H), and 10.09 (s, 1H).

## EXAMPLE 65

5 2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-, 2-(phenylsulfonyl)hydrazide




Step 1. 3-Methoxy-5-phenyl-benzofuran-2-carboxylic acid hydrazide



This was run in accordance with Example 1, Step 2 except 3-methoxy-  
 10 5-phenylbenzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (yield 94%).  
 MS: 283.0 ( $M+1$  for  $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_3$ ); mp: 175.3-177.3°C. HPLC ( $\text{C}_{18}$  column, 1:1  $\text{MeCN}/\text{H}_2\text{O}+0.1\%$  TFA) 99.8%, RT = 1.42 min. Micro: (calc) ( $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_3 \cdot 0.1\text{H}_2\text{O}$ ) C: 67.64, H: 5.04, N: 9.86; (found) C: 67.62, H: 5.05 N: 9.65.  
 15  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  4.20 (s, 3H), 4.52 (s, 2H), 7.35 (t, 1H,  $J = 7.33$  Hz), 7.45 (t, 2H,  $J = 7.33$  Hz), 7.62 (d, 1H,  $J = 8.79$  Hz), 7.69-7.74 (m, 3H), 8.02 (d, 1H,  $J = 1.47$  Hz), and 9.27 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-, 2-(phenylsulfonyl)-hydrazide



20 This was run in accordance with Example 35 except 3-methoxy-5-phenyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-


- 72 -

2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by recrystallization with ethyl acetate/hexanes (53.9% yield).


MS: 423.0 (M+1 for C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 180.7-181.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.4%, RT = 10.39 min. Micro: (calc) (C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S) C: 62.55, H: 4.29, N: 6.63, S: 7.59; (found) C: 62.16, H: 4.05, N: 6.57, S: 7.63; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.07 (s, 3H), 7.35 (t, 1H, J = 8.06 Hz), 7.45 (dd, 2H, J = 1.71 and 7.33 Hz), 7.52 (dd, 2H, J = 1.22 and 7.82 Hz), 7.62 (t, 2H, J = 6.84 Hz), 7.70 (d, 2H, J = 8.55 Hz), 7.76 (dd, 1H, J = 1.95 and 8.79 Hz), 7.82 (d, 2H, J = 7.08 Hz), 8.01 (d, 1H, J = 1.47 Hz), 10.08 (d, 1H, J = 3.2 Hz), and 10.18 (d, 1H, J = 3.2 Hz).

## EXAMPLE 66

2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-, 2-(phenylsulfonyl)hydrazide



Step 1. 3-Ethoxy-5-phenyl-benzofuran-2-carboxylic acid hydrazide




This was run in accordance with Example 1, Step 2 except 3-ethoxy-5-phenylbenzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (yield 44%).

MS: 297.3 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub>); mp: 94.8-96.8°C. HPLC (C18 column, 1:1

MeCN/H<sub>2</sub>O+0.1% TFA) 97.3%, RT = 2.44 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.34 (t, 3H, J = 7.08 Hz), 4.53 (s+q, 4H, J = 6.84 Hz), 7.36 (t, 1H, J = 6.11 Hz), 7.45 (t, 2H, J = 7.33 Hz), 7.62 (d, 1H, J = 8.79 Hz), 7.70-7.73 (m, 3H), 7.95 (d, 1H, J = 1.47 Hz), and 9.22 (s, 1H).

- 73 -

Step 2. 2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-, 2-(phenylsulfonyl)-hydrazide



This was run in accordance with Example 35 except 3-ethoxy-5-phenyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by column chromatography with ethyl acetate/hexanes (42.5% yield).

MS: 437.1 (M+1 for C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 184.3-184.6°C; HPLC (C18 column, 1:1


MeCN/H<sub>2</sub>O+0.1% TFA) 99.9%, RT = 13.67 min. Micro: (calc)

(C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>5</sub>S·0.2H<sub>2</sub>O) C: 62.62, H: 4.66, N: 6.35; (found) C: 62.72, H: 4.52, N:


6.25. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.34 (t, 3H, J = 7.09 Hz), 4.45 (q, 2H, J = 6.83 Hz), 7.45 (t, 1H, J = 7.33 Hz), 7.55 (t, 2H, J = 7.33 Hz), 7.64 (t, 2H, J = 7.81 Hz), 7.72 (t, 2H, J = 5.63 Hz), 7.80 (d, 2H, J = 8.55 Hz), 7.85 (dd, 1H, J = 1.71 and 8.79 Hz), 7.92 (d, 2H, J = 7.08 Hz), 8.05 (d, 1H, J = 1.71 Hz), 10.18 (s, 1H), and 10.18 (s, 1H).

#### EXAMPLE 67

2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-, 2-(phenylsulfonyl)hydrazide



Step 1. 3-Methoxy-7-phenyl-benzofuran-2-carboxylic acid hydrazide




- 74 -

The hydrazide was prepared in accordance with Example 1, Step 2 except that except 3-methoxy-7-phenylbenzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (yield 44%).

MS: 283.2 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>); mp: 177.8-179.13°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 1.42 min. Micro: (calc)

(C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>·0.5H<sub>2</sub>O) C: 65.97, H: 5.19, N: 9.62; (found) C: 66.06, H: 4.93, N: 9.16. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.15 (s, 3H), 4.54 (s, 2H), 7.37-7.52 (m, 2H), 7.50 (dt, 2H, J = 1.71 and 7.33 Hz), 7.64 (dd, 1H, J = 0.98 and 7.33 Hz), 7.78 (dd, 1H, J = 0.98 and 7.33 Hz), 7.90 (dd, 2H, J = 1.47 and 8.55 Hz), and 9.30 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-, 2-(phenylsulfonyl)-hydrazide





This was run in accordance with Example 35 except 3-methoxy-7-phenyl benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by recrystallization with ethyl acetate/hexanes (53.9% yield).

MS: 423.0 (M+1 for C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 186.7-187.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.7%, RT = 10.04 min. Micro: (calc) (C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S) C:

62.55, H: 4.29, N: 6.63, S: 7.59; (found) C: 62.09, H: 4.34, N: 6.58, S: 7.72. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.00 (s, 3H), 7.38-7.41 (m, 2H), 7.41-7.57 (m, 4H), 7.60 (d, 1H, J = 7.57 Hz), 7.67 (dd, 1H, J = 1.23 and 7.58 Hz), 7.77-7.83 (m, 3H), 7.87 (dd, 1H, J = 1.22 and 7.08 Hz), 10.08 (d, 1H, J = 2.8 Hz), and 10.17 (d, 1H, J = 2.8 Hz).


- 75 -

## EXAMPLE 68

Naphtho[2,3-*b*]furan-2-carboxylic acid, 3-methoxy-, 2-(phenylsulfonyl)hydrazideStep 1. 3-Methoxy-naphtho[2,3-*b*]furan-2-carboxylic acid hydrazide

The hydrazide was prepared in accordance with Example 1, Step 2 except that 3-methoxy-naphtho[2,3-*b*]furan-2-carboxylic acid, which was prepared in accordance with the methods of Einhorn et al., *J. Het. Chem.*, 1985:1243, 1247, was used instead of dibenzofuran-2-carboxylic acid (yield 90%).

MS: 257.1 (M+1 for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.4 (t, 1H, J = 3.66 Hz), 3.59 (s, 1H), 4.35 (s, 3H), 5.90 (br., 1H), 7.37-7.46 (m, 2H), 7.85 (s, 2H), 7.88 (d, 1H, J = 9.59 Hz), and 8.21 (s, 1H).

Step 2. Naphtho[2,3-*b*]furan-2-carboxylic acid, 3-methoxy-, 2-(phenylsulfonyl)-hydrazide


This was run in accordance with Example 35 except 3-methoxynaphthofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by chromatography with ethyl acetate/hexanes (38% yield).

MS: 397.0 (M+1 for C<sub>20</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S); mp: 205.7-207.1°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%, RT = 7.61 min. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 4.09 (s, 3H), 7.43-7.49 (m, 1H), 7.53 (t, 3H, J = 7.32 Hz), 7.61 (t, 1H, J = 7.81 Hz), 7.83 (d, 2H, J = 7.33 Hz), 8.0-8.07 (m, 3H), 8.45 (s, 1H), 10.12 (s, 1H), and 10.34 (s, 1H).


- 76 -

## EXAMPLE 69

2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl) sulfonyl hydrazide



Step 1. 5,7-Dichloro-benzofuran-2-carboxylic acid hydrazide



5


The hydrazide was prepared in accordance with Example 1, Step 2 using 5,7-dichloro-benzofuran-2-carboxylic acid that was made according to the procedure in Kurdakar R., *Proc.-Indian Acad. Sci. Sect A*, 1963;58:336, 339 instead of dibenzofuran-2-carboxylic acid (yield 71%).

10

MS: 244.9 (M+1 for  $C_9H_6N_2O_2Cl_2$ ); mp: 237.8-240.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.0%, RT = 2.2 min. Micro: (calc) ( $C_9H_6N_2O_2Cl_2 \cdot 0.65H_2O$ ) C: 42.10, H: 2.87, N: 10.91, Cl: 27.61; (found) C: 42.09, H: 2.50, N: 10.641, Cl: 27.21. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.61 (s, 2H), 7.54 (s, 1H), 7.68 (d, 1H, J = 1.95 Hz), 7.85 (d, 1H, J = 1.95 Hz), and 10.15 (s, 1H).

15

Step 2. 2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl)sulfonyl hydrazide



20


This was run in accordance with Example 35 except that 5,7-dichloro-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride. The desired product was purified by recrystallization in ethyl acetate (52.1% yield).

- 77 -

MS: 384.8 (M+1 for C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 216.4-218.1°C; Micro: (calc) for (C<sub>15</sub>H<sub>10</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 46.77, H: 2.62, N: 7.27, S: 8.32, Cl: 18.41; (found) C: 46.69, H: 2.95, N: 7.14, S: 8.12, Cl: 18.45; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.97%, RT = 8.10 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.53 (t, 2H, J = 7.81 Hz), 7.51-7.61 (m, 1H), 7.66 (s, 1H), 7.72 (d, 1H, J = 1.96 Hz), 7.81 (d, 2H, J = 7.33 Hz), 7.72 (d, 1H, J = 1.96 Hz), 10.27 (d, 1H, J = 2.40 Hz), and 11.10 (d, 1H, J = 2.40 Hz).

## EXAMPLE 70

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide




This was run in accordance with Example 69 except that 2-trifluoromethyl phenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (61.7% yield).

MS: 452.9 (M+1 for C<sub>16</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>3</sub>Cl<sub>2</sub>); mp: 245.6-247.6°C; Micro: (calc) (C<sub>16</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SF<sub>3</sub>Cl<sub>2</sub>) C: 42.40, H: 2.00, N: 6.18, S: 7.07, F: 12.58; (found) C: 42.79, H: 1.99, N: 6.04, S: 6.87, F: 12.31; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.5%, RT = 13.50 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.65 (s, 1H), 7.72 (d, 1H, J = 1.96 Hz), 7.78-7.84 (m, 2H), 7.88 (d, 1H, J = 1.96 Hz), 7.96 (d, 1H, J = 6.60 Hz), 8.16 (d, 1H, J = 6.35 Hz), 10.27 (s, 1H), and 11.18 (s, 1H).

- 78 -

## EXAMPLE 71

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-methylphenyl)sulfonyl]hydrazide



This was run in accordance with Example 69 except that 2-methyl phenyl  
5 sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product  
was purified by recrystallizing in ethyl acetate/hexanes and then chromatographing in  
the same solvent system (28.7% yield).

MS: 398.9 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 212-214°C; Micro: (calc)

(C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 48.13, H: 3.03, N:7.02, S: 8.03; (found) C: 48.08, H: 2.78, N:


10 6.87, S: 8.14; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.9%, RT = 11.86

min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.68 (s, 3H), 7.26 (t, 1H, J = 7.58 Hz), 7.37 (d, 1H,

J = 7.33 Hz), 7.48 (t, 1H, J = 7.33 Hz), 7.63 (s, 1H), 7.71 (d, 1H, J = 1.95 Hz), 7.80 (d,  
1H, J = 7.09 Hz), 7.86 (d, 1H, J = 1.71 Hz), 10.19 (s, 1H), and 11.03 (s, 1H).

## EXAMPLE 72

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-methylphenyl)sulfonyl]-hydrazide



This was run in accordance with Example 69, except that 3-methyl phenyl  
sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product  
was purified by recrystallizing in ethyl alcohol (34.1% yield).

20 MS: 398.9 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); mp: 230.4-231.1°C; Micro: (calc)

(C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>) C: 48.13, H: 3.03, N: 7.02, S:8.03; (found) C: 48.10, H: 2.87, N:

6.96, S: 8.15; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.6%,

RT = 10.45 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.32 (s, 3H), 7.38-7.45 (m, 2H), 7.59 (d, 1H,


J = 7.57 Hz), 7.63 (s, 1H), 7.66 (s, 1H), 7.72 (d, 1H, J = 1.96 Hz), 7.80 (d, 1H,

25 J = 1.71 Hz), 10.23 (s, 1H), and 11.07 (s, 1H).

- 79 -

## EXAMPLE 73

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-bromophenyl)sulfonyl]-hydrazide




This was run in accordance with Example 69 except that 2-bromophenyl  
 5 sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product  
 was purified by column chromatography eluting with ethyl acetate/hexanes 1:2 (26.9%  
 yield).

MS: 462.8 (M+1 for  $C_{15}H_9N_2O_4SCl_2Br$ ); mp: 196.2-196.6°C. HPLC (C18 column,  
 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 14.32 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ  
 10 7.56-7.64 (m, 2H), 7.75 (s, 1H), 7.80 (d, 1H, J = 1.96 Hz), 7.91 (d, 1H, J = 6.84 Hz),  
 7.96 (d, 1H, J = 1.71 Hz), 8.09 (dd, 1H, J = 6.10 and 1.95 Hz), 10.41 (s, 1H), and  
 11.16 (s, 1H).

## EXAMPLE 74

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-bromophenyl)sulfonyl]-hydrazide




This was run in accordance with Example 69 except that 3-bromophenyl  
 sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product  
 was purified by recrystallizing in ethyl acetate (35.2% yield).

MS: 462.8 (M+1 for  $C_{15}H_9N_2O_4SCl_2Br$ ); mp: 248.5-248.8°C; Micro: (calc)  
 20 ( $C_{15}H_9N_2O_4SCl_2Br$ ) C: 48.13, H: 3.03, N: 7.02, S: 8.03; (found) C: 48.10, H: 2.87,  
 N: 6.96, S: 8.15; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.5%,  
 RT = 14.78 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.49 (t, 1H, J = 8.06 Hz), 7.67 (s, 1H), 7.73  
 (d, 1H, J = 1.95 Hz), 7.78 (dd, 1H, J = 6.83 and 0.98 Hz), 7.84 (dd, 1H, J = 0.97 and  
 1.95 Hz), 7.89 (d, 1H, J = 1.96 Hz), 7.94 (s, 1H), 10.50 (d, 1H, J = 2.40 Hz), and 11.17  
 25 (d, 1H, J = 2.40 Hz).

- 80 -

## EXAMPLE 75

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-chlorophenyl)sulfonyl]-hydrazide



This was run in accordance with Example 69 except that 3-chlorophenyl  
 5 sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product  
 was purified by recrystallizing in ethyl acetate (62.2% yield).


MS: 420.9 (M+1 for C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>3</sub>); mp: 245.6-245.7°C; Micro: (calc)

(C<sub>15</sub>H<sub>9</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>3</sub>) C: 42.93, H: 2.16, N: 6.68, S: 7.64, Cl: 25.34; (found) C: 42.96,  
 H: 2.10, N: 6.75, S: 7.73, Cl: 25.08; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA)

10 92.7%, RT = 13.64 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.56, (t, 1H, J = 8.06 Hz), 7.67 (s,  
 1H), 7.71-7.77 (m, 3H), 7.82 (d, 1H, J = 1.71 Hz), 7.89 (d, 1H, J = 1.95 Hz), 10.51 (d,  
 1H, J = 2.40 Hz), and 11.17 (d, 1H, J = 2.40 Hz).

## EXAMPLE 76

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethoxy)phenyl]-  
 15 sulfonyl]hydrazide



This was run in accordance with Example 69 except that  
 2-trifluoromethoxyphenyl sulfonyl chloride was used instead of benzene sulfonyl  
 chloride. The desired product was purified by column chromatography eluting with  
 20 1:2 ethyl acetate/hexanes (33.5% yield).

MS: 468.9 (M+1 for C<sub>16</sub>H<sub>9</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>Cl<sub>2</sub>); mp: 234.5-236.1°C; Micro: (calc)


(C<sub>16</sub>H<sub>9</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>Cl<sub>2</sub>) C: 40.99, H: 1.93, N: 5.97, S: 6.83; (found) C: 41.05, H: 1.88,  
 N: 5.81, S: 6.38; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.1%,  
 RT = 14.82 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.47 (t, 1H, J = 7.81 Hz), 7.53 (d, 1H,

- 81 -


J = 8.31 Hz), 7.65 (s, 1H), 7.72 (d, 1H, J = 1.95 Hz), 7.76 (dd, 1H, J = 1.46 and 8.06 Hz), 7.89 (d, 1H, J = 1.71 Hz), 7.93 (dd, 1H, J = 1.71 and 7.81 Hz), 10.42 (s, 1H), and 11.15 (s, 1H).

## EXAMPLE 77

## 5 Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)hydrazide




Step 1. 5-Methyl-benzofuran-2-carboxylic acid, hydrazide



This was run in accordance with Example 1, Step 2 using 5-methyl-  
 10 benzofuran-2-carboxylic acid that was prepared in accordance to Pappalardo, *Boll. Sci. Fac. Ind. Bologna* 1952:168, instead of dibenzofuran-2-carboxylic acid (95.5% yield). MS: 190.9 (M+1 for C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>); mp: 148.3-150.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.4%, RT = 1.77 min. Micro: (calc) (C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>·0.15H<sub>2</sub>O) C: 62.26, H: 5.38, N: 14.52; (found) C: 62.26, H: 5.38, N:  
 15 14.52. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.37 (s, 3H), 4.41 (d, 2H, J = 3.66 Hz), 7.21 (dd, 1H, J = 8.55 and 1.47 Hz), 7.39 (d, 1H, J = 0.73 Hz), 7.46 (d, 1H, J = 9.28 Hz), 7.49 (s, 1H), and 9.95 (s, 1H).

Step 2. Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)hydrazide




20 This was run in accordance with Example 35 using 5-methylbenzofuran-2-carboxylic acid hydrazide instead of dibenzofuran-2-carboxylic acid hydrazide was used, and benzene sulfonylchloride instead of 3-fluorophenyl sulfonyl chloride was used. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (63.5% yield).

- 82 -


MS: 331.0 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 183.3-183.6°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 5.10 min. Micro: (calc) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S) C: 58.17, H: 4.27, N: 8.48, S: 9.71; (found) C: 58.09, H: 4.30, N: 8.21, S: 9.65. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.55 and 1.71 Hz), 7.47-7.60 (m, 5H), 7.61 (t, 1H, J = 2.20 Hz), 7.80 (dd, 2H, J = 7.08 and 1.47 Hz), 10.16 (s, 1H), and 10.88 (s, 1H).

## EXAMPLE 78

2-Benzofurancarboxylic acid, 5-chloro-3-methoxy, 2-(phenylsulfonyl)hydrazide




Step 1. 5-Chloro-3-methoxy-2-carboxylic acid hydrazide



This was run in accordance with Example 1, Step 2 except 5-chloro-3-methoxy-benzofuran-2-carboxylic acid, which was prepared in accordance with the methods of Perkin, *J. Chem. Soc.*, 1951:101-103, was used instead of dibenzofuran-2-carboxylic acid (76% yield).

MS: 241.0.9 (M+1 for C<sub>10</sub>H<sub>9</sub>N<sub>2</sub>O<sub>3</sub>Cl); mp: 205.6-206.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.7%, RT = 2.12 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.12 (s, 3H), 4.52 (s, 2H), 7.46 (dd, 1H, J = 8.79 and 2.19 Hz), 7.58 (d, 1H, J = 8.79 Hz), 7.91 (d, 1H, J = 2.20 Hz), and 9.31 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 5-chloro-3-methoxy-, 2-(phenylsulfonyl)-hydrazide




- 83 -

This was run in accordance with Example 35 except 5-chloro-3-methoxy-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide was used, and benzene sulfonylchloride instead of 3-fluorophenyl sulfonyl chloride was used (28.5% yield).

MS: 381.0 (M+1 for  $C_{16}H_{13}N_2O_5SCl$ ); mp: 242.2-244.2°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.0%, RT = 6.38 min. Micro: (calc) ( $C_{16}H_{13}N_2O_5SCl \cdot 0.3H_2O$ ) C: 49.64, H: 3.54, N: 7.24, S: 8.28; (found) C: 49.78, H: 3.23, N: 7.23, S: 8.56. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 4.01 (s, 3H), 7.48-7.54 (m, 3H), 7.58-7.64 (m, 2H), 7.81 (d, 2H, J = 7.57 Hz), 7.91 (d, 1H, J = 1.96 Hz), 10.09 (s, 1H), and 10.23 (s, 1H).

## EXAMPLE 79

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethyl)phenyl]sulfonyl]-hydrazide




This was run in accordance with Example 77 except 2-trifluoromethyl phenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (62.5% yield).

MS: 398.9 (M+1 for  $C_{17}H_{13}N_2O_4SF_3$ ); mp: 194.0-195.4°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.8%, RT = 8.34 min. Micro: (calc) ( $C_{17}H_{13}N_2O_4SF_3$ ) C: 51.26, H: 3.29, N: 7.03, S: 8.05, F: 14.31; (found) C: 51.24, H: 3.40, N: 6.87, S: 7.93, F: 13.87. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.30 and 1.71 Hz), 7.47-7.51 (m, 3H), 7.76-7.83 (m, 2H), 7.95 (dd, 2H, J = 7.08 and 1.96 Hz), 8.15 (dd, 2H, J = 6.84 and 2.20 Hz), 10.15 (s, 1H), and 10.96 (s, 1H).

- 84 -

## EXAMPLE 80


2-Benzofurancarboxylic acid, 5-methyl-, 2-([3-methylphenyl]sulfonyl)hydrazide



This was run in accordance with Example 77 except 3-methylphenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (64.6% yield). MS: 345.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 184.1-186.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.28%, RT = 5.91 min.; Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S) C: 59.29, H: 4.69, N: 8.14, S: 9.29; (found) C: 59.15, H: 4.67, N: 8.13, S: 9.56. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.31 (s, 3H), 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.31 and 1.95 Hz), 7.37-7.43 (m, 2H), 7.48-7.51 (m, 3H), 7.59 (d, 1H, J = 7.33 Hz), 7.62 (s, 1H), 10.11 (s, 1H), and 10.84 (s, 1H).

## EXAMPLE 81

2-Benzofurancarboxylic acid, 5-methyl-, 2-[2-(methylphenyl)sulfonyl]hydrazide




This was run in accordance with Example 77 except 2-methylphenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (57.0% yield). MS: 345.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 174.5-174.7°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.6%, RT = 6.68 min. Micro: (calc) (C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S) C: 59.29, H: 4.69, N: 8.14, S: 9.29; (found) C: 59.21, H: 4.51, N: 8.05, S: 9.38. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 2.67 (s, 3H), 7.25 (t, 2H, J = 7.57 Hz), 7.35 (d, 1H, J = 7.32 Hz), 7.45-7.49 (m, 4H), 7.79 (dd, 1H, J = 7.82 and 1.22 Hz), 10.07 (s, 1H), and 10.79 (s, 1H).

- 85 -

## EXAMPLE 82

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chloro-2-methylphenyl)sulfonyl]-hydrazide




This was run in accordance with Example 77 except 3-chloro-2-methylphenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by column chromatography eluting with ethyl acetate/hexanes (40.2% yield).

MS: 378.9 (M+1 for  $C_{17}H_{15}N_2O_4SCl$ ); mp: 182.4-184.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.2%, RT = 11.05 min. Micro: (calc) ( $C_{17}H_{15}N_2O_4SCl$ ) C: 53.90, H: 3.99, N: 7.39, S: 8.46, Cl: 9.36; (found) C: 54.13, H: 4.03, N: 7.27, S: 8.41, Cl: 9.31. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 2.71 (s, 3H), 7.25 (dd, 1H, J = 8.55 and 1.71 Hz), 7.29 (t, 1H, J = 8.06 Hz), 7.46-7.50 (m, 3H), 7.69 (d, 1H, J = 8.06 Hz), 7.81 (dd, 1H, J = 8.06 and 0.98 Hz), 10.34 (s, 1H), and 10.85 (s, 1H).

## EXAMPLE 83

2-Benzofurancarboxylic acid, 5-methyl-, 2-[3-(bromophenyl)sulfonyl]hydrazide



This was run in accordance with Example 77 except 3-bromophenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by recrystallizing in ethyl acetate/hexanes (58.7% yield).


MS: 408.9 (M+1 for  $C_{16}H_{13}N_2O_4SBr$ ); mp: 207.2-209.9°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.8%, RT = 8.77 min. Micro: (calc) ( $C_{16}H_{13}N_2O_4SBr$ ) C: 46.96, H: 3.20, N: 6.84, S: 7.83, Br: 19.52; (found) C: 46.99, H: 3.05, N: 6.77, S: 8.30, Br: 20.24. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.79 and 1.71

- 86 -

Hz), 7.44 (d, 1H, J = 7.58 Hz), 7.46-7.51 (m, 4H), 7.74 (dq, 1H, J = 7.57 and 1.71 Hz), 7.92 (dd, 1H, J = 7.81 and 1.71 Hz), 10.30 (s, 1H), and 10.92 (s, 1H).

## EXAMPLE 84

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chlorophenyl)sulfonyl]hydrazide




This was run in accordance with Example 77 except 3-chlorophenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by chromatographing with ethyl acetate/hexanes (75.3% yield).

MS: 365.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl); mp: 186.1-186.3°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.7%, RT = 8.08 min. Micro: (calc) (C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl) C: 52.68, H: 3.59, N: 7.68, S: 8.79, Cl: 9.72; (found) C: 52.64, H: 3.60, N: 7.59, S: 8.71, Cl: 9.89. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.54 and 1.22 Hz), 7.48-7.57 (m, 3H), 7.70-7.75 (m, 2H), 7.81 (s, 1H), 10.40 (s, 1H), and 10.96 (s, 1H).

## EXAMPLE 85

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethoxy)phenyl]-sulfonyl]hydrazide




This was run in accordance with Example 77 except 2-trifluoromethoxy phenyl sulfonyl chloride was used instead of benzene sulfonyl chloride. The desired product was purified by chromatographing with ethyl acetate/hexanes (72.5% yield). MS: 415.0 (M+1 for C<sub>17</sub>H<sub>17</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>); mp: 226.7-227.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.8%, RT = 8.83 min. Micro: (calc)

- 87 -


(C<sub>17</sub>H<sub>17</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>3</sub>) C: 49.28, H: 3.16, N: 6.76, S: 7.74, F: 13.75; (found) C: 49.18, H: 3.10, N: 6.67, S: 7.84, F: 14.00. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.36 (s, 3H), 7.25 (dd, 1H, J = 8.55 and 1.71 Hz), 7.44 (dd, 1H, J = 8.55 and 0.97 Hz), 7.47-7.51 (m, 4H), 7.74 (dt, 1H, J = 8.30 and 1.71 Hz), 7.92 (dd, 1H, J = 7.82 and 1.47 Hz), 10.30 (s, 1H), and 10.92 (s, 1H).

## EXAMPLE 86

2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)hydrazide




Step 1. 7-Methylbenzofuran-2-carboxylic acid hydrazide



This was run in accordance with Example 1, Step 2 except 7-methylbenzofuran-2-carboxylic acid that was prepared in accordance to Pappalardo, *Boll. Sci. Fac. Ind. Bologna*, 1952:168, was used instead of dibenzofuran-2-carboxylic acid (60% yield).

MS: 190.9 (M+1 for C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>); mp: 171.6-173.4°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.2%, RT = 1.79 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.48, (s, 3H), 4.55 (s, 2H), 7.15-7.22 (m, 2H), 7.44 (s, 1H), 7.51 (dd, 1H, J = 6.84 and 0.98 Hz), and 9.98 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)hydrazide



This was run in accordance with Example 35 except 7-methylbenzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride.

- 88 -


The desired product was purified by recrystallizing in ethyl acetate/hexanes (71% yield).

MS: 331.0 (M+1 for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 217.5-218.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 97.8%, RT = 5.14 min. Micro: (calc) (C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S) C:


5 58.17, H: 4.27, N: 8.48, S: 9.71; (found) C: 57.87, H: 4.26, N: 8.37, S: 9.70. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.48 (s, 3H), 7.18 (t, 1H, J = 7.57 Hz), 7.24 (d, 1H, J = 7.09 Hz), 7.50-7.54 (m, 4H), 7.62 (t, 1H, J = 7.57 Hz), 7.82 (d, 2H, J = 7.33 Hz), 10.19 (d, 1H, J = 2.4 Hz), and 10.91 (d, 1H, J = 2.4 Hz).

## EXAMPLE 87

10 2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)hydrazide



Step 1. 4-Methylbenzofuran-2-carboxylic acid hydrazide




15 This was run in accordance with Example 1, Step 2 except 4-methylbenzofuran-2-carboxylic acid that was prepared in accordance to Pappalardo, *Boll. Sci. Fac. Ind. Bologna*, 1952:168, was used instead of dibenzofuran-2-carboxylic acid (60% yield).

MS: 190.9 (M+1 for C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>); mp: 175.4-177.0°C. HPLC (C18 column, 1:1

20 MeCN/H<sub>2</sub>O+0.1% TFA) 95.4%, RT = 1.76 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.46 (s, 3H), 4.52 (d, 2H, J = 3.91 Hz), 7.07 (d, 1H, J = 7.33 Hz), 7.29 (t, 1H, J = 8.30 Hz), 7.39 (d, 1H, J = 8.30 Hz), 7.54 (d, 1H, J = 0.97 Hz), and 9.95 (s, 1H).

- 89 -

Step 2. 2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)hydrazide




This was run in accordance with Example 35 except 4-methyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride (91% yield).


MS: 331.1 (M+1 for  $C_{16}H_{14}N_2O_4S$ ); mp: 217.0-218.0°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.3%, RT = 4.80 min. Micro: (calc) ( $C_{16}H_{14}N_2O_4S$ ) C: 58.17, H: 4.27, N: 8.48, S: 9.71; (found) C: 57.73, H: 4.22, N: 8.31, S: 9.71. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.46 (s, 3H), 7.10 (d, 1H, J = 7.32 Hz), 7.32 (t, 1H, J = 8.30 Hz), 7.39 (d, 1H, J = 8.30 Hz), 7.49-7.53 (m, 2H), 7.60 (dd, 1H, J = 5.62 and 1.22 Hz), 7.61 (s, 1H), 7.80 (d, 2H, J = 7.32 Hz), 10.16 (s, 1H), and 10.87 (s, 1H).

## EXAMPLE 88

2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)hydrazide



Step 1. 3-Methylbenzofuran-2-carboxylic acid hydrazide




This was run in accordance with Example 1, Step 2 except 3-methyl-benzofuran-2-carboxylic acid was used instead of dibenzofuran-2-carboxylic acid (74% yield).

MS: 191.0 (M+1 for  $C_{10}H_{10}N_2O_2$ ); mp: 136-137°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 95.3%, RT = 1.87 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.52 (s, 3H),

- 90 -

4.48 (s, 2H), 7.29 (dt, 1H,  $J = 7.82$  and  $0.74$  Hz), 7.39-7.43 (m, 1H), 7.51 (d, 1H,  $J = 8.31$  Hz), 7.68 (d, 1H,  $J = 7.82$  Hz), and 9.79 (s, 1H).

Step 2. 2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)hydrazide




5 This was run in accordance with Example 35 except 3-methyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenyl sulfonyl chloride was used instead of 3-fluorophenyl chloride (68.5% yield).  
 MS: 331.1 ( $M+1$  for  $C_{16}H_{14}N_2O_4S$ ); mp: 221.6-222.2°C. HPLC (C18 column, 1:1  
 10 MeCN/ $H_2O$ +0.1% TFA) 99.4%, RT = 5.50 min. Micro: (calc) ( $C_{16}H_{14}N_2O_4S$ ) C: 58.17, H: 4.27, N: 8.48, S: 9.71; (found) C: 58.31, H: 4.22, N: 8.55, S: 9.73.  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  2.35 (s, 3H), 7.31 (t, 1H,  $J = 8.06$  Hz), 7.46 (t, 1H,  $J = 8.30$  Hz), 7.50-7.60 (m, 3H), 7.61-7.63 (m, 1H), 7.69 (d, 1H,  $J = 7.81$  Hz), 7.81 (d, 2H,  $J = 7.32$  Hz), 10.07 (s, 1H), and 10.70 (s, 1H).


15

## EXAMPLE 89

2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)hydrazide



Step 1. 3,5-di-Methylbenzofuran-2-carboxylic acid hydrazide




20

This was run in accordance with Example 1, Step 2 except 3,5-dimethyl-benzofuran-2-carboxylic acid, which was prepared in accordance to Dey, *J. Chem. Soc.*, 1945:1646, was used instead of dibenzofuran-2-carboxylic acid (40% yield).

- 91 -

MS: 190.9 (M+1 for C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>); mp: 142.7-145.1°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 95.1%, RT = 1.82 min. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.37 (s, 3H), 2.47 (s, 3H), 4.47 (s, 2H), 7.21 (dd, 1H, J = 8.54 and 1.46 Hz), 7.38 (d, 1H, J = 8.30 Hz), 7.45 (d, 1H, J = 0.73 Hz), and 9.74 (s, 1H).

5 Step 2. 2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)hydrazide




This was run in accordance with Example 35 except 3,5-dimethyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenylsulfonyl chloride was used instead of 3-fluorophenyl chloride (47.3% yield).

MS: 345.1 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 225.9-227.1°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 99.0%, RT = 7.20 min. Micro: (calc)


(C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S·0.6H<sub>2</sub>O) C: 57.34, H: 4.87, N: 7.87, S: 9.00; (found) C: 57.46, H: 4.48, N: 7.90, S: 9.37. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.31 (s, 3H), 2.37 (s, 3H), 7.25 (dd, 1H, J = 1.47 and 8.80 Hz), 7.42 (d, 1H, J = 8.55 Hz), 7.45 (s, 1H), 7.51 (t, 2H, J = 7.82 Hz), 7.58-7.62 (m, 1H), 7.80 (d, 2H, J = 7.08 Hz), 10.4 (s, 1H), and 10.64 (s, 1H).

#### EXAMPLE 90

2-Benzofurancarboxylic acid, 5-chloro-3-methyl-, 2-(phenylsulfonyl)hydrazide



20 Step 1. 5-Chloro-3-methyl-benzofuran-2-carboxylic acid hydrazide




This was run in accordance with Example 1, Step 2 except 5-chloro-3-methyl-benzofuran-2-carboxylic acid, which was prepared in accordance to Suzuki et al.,

- 92 -

*Bull. Chem. Soc. Jpn.*, 1983:2762-2767, was used instead of dibenzofuran-2-carboxylic acid (40% yield).

MS: 225.1 (M+1 for C<sub>10</sub>H<sub>9</sub>ClN<sub>2</sub>O<sub>2</sub>); mp: 214.3-214.7°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.4%, RT = 1.98 min.

5 Step 2. 2-Benzofurancarboxylic acid, 5-chloro-3-methyl, 2-(phenylsulfonyl)-hydrazide



This was run in accordance with Example 35 except 5-chloro-3-methyl-benzofuran-2-carboxylic acid hydrazide was used instead of dibenzofuran-2-carboxylic acid hydrazide, and phenylsulfonyl chloride was used instead of 3-fluorophenyl chloride (43.2% yield).


10 MS: 365.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl); mp: 246.8-247.8°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 98.6%, RT = 8.24 min. Micro: (calc)

(C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>SCl·0.5H<sub>2</sub>O) C: 51.31, H: 3.77, N: 7.48, S: 8.56; (found) C: 51.46, H:

3.44, N: 8.16, S: 8.86; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.30 (s, 3H), 7.45 (dd, 1H, J = 2.20 and 8.80 Hz), 7.42 (d, 1H, J = 8.55 Hz), 7.51 (t, 2H, J = 7.32 Hz), 7.56-7.61 (m, 2H), 7.78-7.80 (m, 32H), 10.09 (d, 1H, J = 2.8 Hz), and 10.75 (d, 1H, J = 2.8 Hz).

#### EXAMPLE 91

2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide



20 A solution of Example 34 (0.11 g, 0.305 mmol) in THF (20 mL) was treated with H<sub>2</sub> at 40 psi in the presence of Raney Ni (trace) in a Parr shaker. The catalyst was removed by filtration, the solvents removed, and the product was isolated by column chromatography eluting with ethyl acetate (50% yield).

MS: 332.0 (M+1 for C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 206.9-207.7°C. HPLC (C18 column, 1:1

25 MeCN/H<sub>2</sub>O+0.1% TFA) 99.46%, RT = 1.54 min. Micro: (calc) (C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>4</sub>S) C:


- 93 -

54.37, H: 3.95, N: 12.68, S: 9.68; (found) C: 54.18, H: 4.14, N: 12.44, S: 9.56.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  4.99 (s, 2H), 6.71, (d, 2H,  $J = 2.20$  Hz), 7.25 (d, 1H,  $J = 9.52$  Hz), 7.30 (s, 1H), 7.51 (t, 2H,  $J = 7.82$  Hz), 7.61 (t, 1H,  $J = 7.33$  Hz), 7.79 (d, 2H,  $J = 7.57$  Hz), 10.10 (s, 1H), and 10.73 (s, 1H).

5

## EXAMPLE 92

2-Dibenzofurancarboxylic acid, 2-[(2-fluorophenyl)sulfonyl]hydrazide



10

15


20

Synthesis of 2-Dibenzofurancarboxylic acid 2-[(2-fluorophenyl)sulfonyl]-hydrazide: A solution of dibenzofuran-2-carboxylic acid hydrazide (0.3 M) in pyridine (0.513 mL) was treated with a solution of the 2-fluorobenzenesulfonyl chloride (1.0 M) in  $\text{CH}_2\text{Cl}_2$  (0.225 mL). The reaction was placed in a Bohdan Miniblock apparatus and shaken at  $45^\circ\text{C}$  for 16 hours. The reaction was cooled to room temperature, treated with polymer-supported polyamine quench resin (Aldrich, 100 mg), and shaken for 16 hours. The solution was filtered and concentrated. The residue was purified by preparative HPLC on a Phenomenex Develofil  $28 \times 100$  mm C18 column eluting with a gradient of 10% to 100%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  + 3% n-propanol over 6.5 minutes.

MS: 385.4 ( $M+1$  for  $\text{C}_{19}\text{H}_{13}\text{N}_2\text{O}_4\text{SF}$ ); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100%  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.59 min., 100%.

## EXAMPLE 93

2-Dibenzofurancarboxylic acid, 2-(2,6-difluorophenylsulfonyl)hydrazide




- 94 -

Example 93 was synthesized in accordance with the methods of Example 92 except that 2,6-difluorobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 403.4 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>F<sub>2</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.78 min., 100%.

## EXAMPLE 94

2-Dibenzofurancarboxylic acid, 2-[(2-bromophenyl)sulfonyl]hydrazide




Example 94 was synthesized in accordance with the methods of Example 92 except that 2-bromobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 446.3 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Br); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.22 min., 76.0%.

## EXAMPLE 95

2-Dibenzofurancarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide




Example 95 was synthesized in accordance with the methods of Example 92 except that 3-bromobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 445.3 (M + 1 for C<sub>19</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Br); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.59 min., 100%.

- 95 -

## EXAMPLE 96

2-Dibenzofurancarboxylic acid, 2-[(4-methyl-3-nitrophenyl)sulfonyl]hydrazide




Example 96 was synthesized in accordance with the methods of Example 92 except that 4-methyl-3-nitrobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 426.4 (M+1 for C<sub>20</sub>H<sub>15</sub>N<sub>3</sub>O<sub>6</sub>S<sub>1</sub>), HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.21 min., 92.7%.

## EXAMPLE 97

2-Dibenzofurancarboxylic acid, 2-[2-chloro-5-(trifluoromethyl)phenyl]sulfonyl)-hydrazide




Synthesis of dibenzofuran-2-carboxylic acid 2-(2-chloro-5-trifluoromethylphenylsulfonyl)hydrazide: Example 97 was synthesized in accordance with the methods of Example 92 except that 2-chloro-5-trifluoromethylbenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 469.8 (M+1 for C<sub>20</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S<sub>1</sub>Cl F<sub>3</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with a 3 minute run time) RT = 2.33 min., 87.7%.

- 96 -

## EXAMPLE 98

2-Dibenzofurancarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide




Synthesis of 2-dibenzofurancarboxylic acid 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide: Example 98 was synthesized in accordance with the methods of Example 92 except that 2-trifluoromethoxybenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 451.4 (M +1 for C<sub>20</sub>H<sub>13</sub>F<sub>3</sub>N<sub>2</sub>O<sub>5</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.28 min., 94.7%.

## EXAMPLE 99

2-Dibenzofurancarboxylic acid, 2-[(4-bromo-2-trifluoromethoxyphenyl)sulfonyl]hydrazide




Synthesis of 2-dibenzofurancarboxylic acid 2-[(4-bromo-2-trifluoromethoxyphenyl)sulfonyl]hydrazide: Example 99 was synthesized in accordance with the methods of Example 92 except that 4-bromo-2-trifluoromethoxybenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 530.3 (M+1 for C<sub>20</sub>H<sub>12</sub>BrF<sub>3</sub>N<sub>2</sub>O<sub>5</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.42 min., 95.1%.

- 97 -

## EXAMPLE 100

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]hydrazide



Synthesis of 2-dibenzofurancarboxylic acid 2-[(2-nitrophenyl)-


5        sulfonyl]hydrazide: Example 100 was synthesized in accordance with the methods of Example 92 except that 2-nitrobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.

MS: 412.4 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>3</sub>O<sub>6</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with

10        3 minute run time) RT = 3.03 min., 88.3%.

## EXAMPLE 101

2-Dibenzofurancarboxylic acid, 2-[(4-carboxyphenyl)sulfonyl]hydrazide



Synthesis of 2-dibenzofurancarboxylic acid 2-[(4-carboxyphenyl)-

15        sulfonyl]hydrazide: Example 101 was synthesized in accordance with the methods of Example 92 except that 4-carboxybenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride.


MS: 411.4 (M+1 for C<sub>20</sub>H<sub>14</sub>N<sub>3</sub>O<sub>6</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with

20        3 minute run time) RT = 1.85 min., 89.2%.

- 98 -

## EXAMPLE 102

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)sulfonyl]hydrazide




Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)sulfonyl]hydrazide: A solution of 2-(7-methoxyl)-benzofurancarboxylic acid hydrazide (0.3 M) in pyridine (0.513 mL) was treated with a solution of the 3-chlorobenzenesulfonyl chloride (1.0 M) in CH<sub>2</sub>Cl<sub>2</sub> (0.225 mL). The reaction was placed in a Bohdan Miniblock apparatus and shaken at 45°C for 16 hours. The reaction was cooled to room temperature, treated with polymer-supported polyamine quench resin (Aldrich, 100 mg), and shaken for 16 hours. The solution was filtered and concentrated. The residue was purified by preparative HPLC on a Phenomenex Develofil 28 × 100 mm C18 column eluting with a gradient of 10% to 100% CH<sub>3</sub>CN /H<sub>2</sub>O+3% n-propanol over 6.5 minutes.

MS: 380.8 (M+1 for C<sub>16</sub>H<sub>13</sub>ClN<sub>2</sub>O<sub>5</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.67 min., 98.9%.

## EXAMPLE 103

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-fluorophenyl)sulfonyl]hydrazide




Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide: Example 103 was synthesized in accordance with the methods of Example 102 except that 3-fluorobenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

- 99 -

MS: 364.4 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>5</sub>SF); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.47 min., 100%.

## EXAMPLE 104

5 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)sulfonyl]-hydrazide




Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)-sulfonyl]hydrazide. Example 104 was synthesized in accordance with the methods of Example 102 except that 2,3-dichlorobenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

10 MS: 415.3 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>SCl<sub>2</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.79 min., 92.7%.

## EXAMPLE 105

15 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide



Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide: Example 105 was synthesized in accordance with the methods of Example 102 except that 3-chloro-4-fluorobenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.


20

- 100 -

MS: 398.8 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>5</sub>SClF); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.75 min., 100%.

## EXAMPLE 106

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)-sulfonyl]hydrazide




Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)sulfonyl]hydrazide: Example 106 was synthesized in accordance with the methods of Example 102 except that 5-chloro-2-methoxybenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

MS: 410.8 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SCl); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.69 min., 97.3%.

## EXAMPLE 107

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)-sulfonyl]hydrazide




Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)sulfonyl]hydrazide: Example 107 was synthesized in accordance with the methods of Example 102 except that 5-bromo-2-methoxybenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

- 101 -

MS: 455.3 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>SBr); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.74 min., 97.8%.

## EXAMPLE 108

## 5 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)sulfonyl]hydrazide




10 Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)sulfonyl]hydrazide: Example 108 was synthesized in accordance with the methods of Example 102 except that 4-ethylbenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

MS: 374.4 (M+1 for C<sub>18</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.75 min., 86.1%.

## EXAMPLE 109

## 15 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide




20 Synthesis of 2-benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)sulfonyl]hydrazide: Example 109 was synthesized in accordance with the methods of Example 102 except that 3-methoxybenzenesulfonyl chloride was used instead of 3-chlorobenzenesulfonyl chloride.

MS: 376.4 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 3.45 min., 100%.

- 102 -

## EXAMPLE 110

2-Dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide




Synthesis of 2-dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)-  
sulfonyl]hydrazide: Example 110 was synthesized in accordance with the methods of  
Example 92 except that 2,3-dichlorobenzenesulfonyl chloride was used instead of  
2-fluorobenzenesulfonyl chloride.

MS: 436.3 (M+1 for C<sub>19</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>SCl<sub>2</sub>); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
3 minute run time) RT = 4.18 min., 94.5%.

## EXAMPLE 111

2-Naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-  
hydrazide: A solution of 2-naphthalenecarboxylic acid, hydrazide (0.3 M) in pyridine  
(0.513 mL) was treated with a solution of the 3-methylbenzenesulfonyl chloride  
(1.0 M) in CH<sub>2</sub>Cl<sub>2</sub> (0.225 mL). The reaction was placed in a Bohdan Miniblock  
apparatus and shaken at 45°C for 16 hours. The reaction was cooled to room  
temperature, treated with polymer-supported polyamine quench resin (Aldrich,  
100 mg), and shaken for 16 hours. The solution was filtered and concentrated. The  
residue was purified by preparative HPLC on a Phenomenex Develofil 28 × 100 mm  
C18 column eluting with a gradient of 10% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O+3% n-propanol  
over 6.5 minutes.

- 103 -

MS: 341.4 (M+1 for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.18 min., 72.0%.

## EXAMPLE 112

5 2-Naphthalenecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(4-methylphenyl)-sulfonyl]hydrazide: Example 112 was synthesized in accordance with the methods of Example 111 except that 4-methylbenzenesulfonyl chloride was used instead of 3-methylbenzenesulfonyl chloride.

MS: 341.4 (M+1 for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 1.59 min., 100%.

## EXAMPLE 113

15 2-Naphthalenecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide: Example 113 was synthesized in accordance with the methods of Example 111 except that 2-trifluoromethylphenylsulfonyl chloride was used instead of 3-methylbenzenesulfonyl chloride.

MS: 395.4 (M+1 for C<sub>18</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.16 min., 84.3%.

- 104 -

## EXAMPLE 114

2-Naphthalenecarboxylic acid, 2-[[4-(trifluoromethyl)phenyl]sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[[4-(trifluoromethyl)-  
phenyl]sulfonyl]hydrazide: Example 114 was synthesized in accordance with the  
5 methods of Example 111 except that 4-trifluoromethylphenylsulfonyl chloride was  
used instead of 3-methylbenzenesulfonyl chloride.

MS: 395.4 (M+1 for C<sub>18</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
10 3 minute run time) RT = 2.13 min., 97.0%.

## EXAMPLE 115

2-Naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-  
sulfonyl]hydrazide: Example 115 was synthesized in accordance with the methods of  
15 Example 111 except that 3-chloro-4-fluorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 379.9 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>SClF); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
20 3 minute run time) RT = 3.90 min., 97.8%.

- 105 -

## EXAMPLE 116

2-Naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)-  
sulfonyl]hydrazide: Example 116 was synthesized in accordance with the methods of  
Example 111 except that 3-methoxyphenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 357.4 (M+1 for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub>S); HPLC (Waters Alliance 2790 column, solvent  
gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
3 minute run time) RT = 3.60 min., 99.9%.

## EXAMPLE 117

2-Naphthalenecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(3-chlorophenyl)-  
sulfonyl]hydrazide: Example 117 was synthesized in accordance with the methods of  
Example 111 except that 3-chlorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 361.8 (M+1 for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>SCl); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
3 minute run time) RT = 3.69 min., 100%.

- 106 -

## EXAMPLE 118

2-Naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-  
5 hydrazide: Example 118 was synthesized in accordance with the methods of  
Example 111 except that 2-chlorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 361.8 (M+1 for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>SCl); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
10 3 minute run time) RT = 3.67 min., 99.9%.

## EXAMPLE 119

2-Naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-  
15 hydrazide: Example 119 was synthesized in accordance with the methods of  
Example 111 except that 3-fluorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 345.4 (M+1 for C<sub>17</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>SF); HPLC (Waters Alliance 2790 column, solvent  
gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
20 3 minute run time) RT = 3.61 min., 99.9%.

- 107 -

## EXAMPLE 120

2-Naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)-  
5 sulfonyl]hydrazide: Example 120 was synthesized in accordance with the methods of  
Example 111 except that 2,3-dichlorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 396.3 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>SCl<sub>2</sub>); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
10 3 minute run time) RT = 3.95 min., 99.4%.

## EXAMPLE 121

2-Naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]hydrazide




Synthesis of 2-naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]-  
15 hydrazide: Example 121 was synthesized in accordance with the methods of  
Example 111 except that 4-ethanysulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 355.4 (M+1 for C<sub>19</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>S); HPLC (Waters Alliance 2790 column, solvent  
gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
20 3 minute run time) RT = 3.90 min., 100%.

- 108 -

## EXAMPLE 122

3-Quinolinecarboxylic acid, 2-[(4-pentylphenyl)sulfonyl]hydrazide




3-Quinolinecarboxylic acid hydrazide (1.0 g, 5.34 mmol) was dissolved in THF (60 mL). N,N-diisopropylethylamine (3.72 mL, 21.36 mmol) was added, and the reaction was cooled to 0°C. 4-Pentylphenylsulfonyl chloride (0.67 mL, 5.34 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 3.5 hours of stirring, the red solution was washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and recrystallized from dichloromethane. The desired product was isolated (0.155 g, 8.9%).

MS: 398.5 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.17 min., 100%.

## EXAMPLE 123

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]sulfonyl]hydrazide




Synthesis of 3-quinolinecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]sulfonyl]hydrazide: Example 123 was synthesized in accordance with the methods of Example 122 except that 2-trifluoromethylphenylsulfonyl chloride was used instead of 4-pentylphenylsulfonyl chloride.

MS: 396.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 1.93 min., 100%.

- 109 -

## EXAMPLE 124


3-Quinolinecarboxylic acid, 2-[[2,5-bis(2,2,2-trifluoroethoxyl)phenyl]-sulfonyl]hydrazide



- 5            Synthesis of 3-quinolinecarboxylic acid, [[2,5-bis(2,2,2-trifluoroethoxyl)phenyl]sulfonyl]hydrazide: Example 124 was synthesized in accordance with the methods of Example 122 except that [2,5-bis(2,2,2-trifluoroethoxyl)phenyl]sulfonyl chloride was used instead of 4-pentylphenylsulfonyl chloride.
- 10           MS: 524.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 2.07 min., 100%.

## EXAMPLE 125

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide




- 15           Synthesis of 3-quinolinecarboxylic acid, [(3-methylphenyl)sulfonyl]-hydrazide: Example 125 was synthesized in accordance with the methods of Example 122 except that (3-methylphenylsulfonyl chloride was used instead of 4-pentylphenylsulfonyl chloride.
- 20           MS: 342.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column, solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with 3 minute run time) RT = 1.83 min., 100%.

- 110 -

## EXAMPLE 126

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]sulfonyl]hydrazide




Synthesis of 3-quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]-  
sulfonyl]hydrazide: Example 126 was synthesized in accordance with the methods of  
Example 122 except that 2-(2-trifluoromethoxy)-phenylsulfonyl chloride was used  
instead of 4-pentylphenylsulfonyl chloride.

MS: 412.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
3 minute run time) RT = 1.94 min., 99.5%.

## EXAMPLE 127

3-Quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]hydrazide




Synthesis of 3-quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-  
hydrazide: Example 127 was synthesized in accordance with the methods of  
Example 122 except that (4-methylphenyl)sulfonyl chloride was used instead of  
4-pentylphenylsulfonyl chloride.

MS: 342.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>); HPLC (Waters Alliance 2790 column,  
solvent gradient of 60% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O + 0.1% formic acid over 1 minute with  
3 minute run time) RT = 1.98 min., 98.3%.

- 111 -

## EXAMPLE 128

6-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-

5 hydrazide: To a suspension of 6-quinolinecarboxylic acid hydrazide (0.1 g, 0.53 mmol) in pyridine (7 mL). Then the 3-methylbenzenesulfonyl chloride (0.107 g, 0.55 mmol) was added, and the reaction was stirred at room temperature overnight. The solvent was removed, and the desired product was purified by recrystallization in ethyl acetate.

10 MS: 342.4 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>3</sub>S); mp: 234.0-234.6°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 96.6%, RT = 1.77 min., <sup>1</sup>H NMR (400 MHz DMSO) δ 2.29(s, 3H), 7.40(m, 2H), 7.67(m, 2H), 7.86(t, 1H, J=6.5 Hz), 8.12(d, 1H, J=8.5 Hz), 8.22(d, 1H, J=8.8 Hz), 8.53(s, 1H), 8.84(d, 1H, J=8.0 Hz), 9.16(s, 1H), 10.14(s, 1H), 11.01(s, 1H).

## EXAMPLE 129

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]hydrazide



Synthesis of 3-quinolinecarboxylic acid, 2-[(3-methylphenyl)-


20 sulfonyl]hydrazide: Example 129 was synthesized in accordance with the methods of Example 128 except that 3-quinolinecarboxylic acid hydrazide was used instead of 6-quinolinecarboxylic acid hydrazide.

MS: 342.4 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>3</sub>S; mp: 238.6-239.4°C. HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 95.9%, RT = 2.43 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ 2.30(s, 3H), 7.40(m, 2H), 7.65(m, 2H), 7.79(t, 1H, J=7.7Hz), 7.99(t, 1H, J=7.9 Hz), 8.18(m, 2H), 8.91(s, 1H), 9.22(s, 1H), 10.22(s, 1H), 11.14(s, 1H).

- 112 -

## EXAMPLE 130

6-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide



Synthesis of 6-quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-


5 hydrazide: Example 130 was synthesized in accordance with the methods of Example 128 except that 3-chlorophenylsulfonyl chloride was used instead of 3-methylbenzenesulfonyl chloride.

MS: 362.8 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SCl); mp: 238.4-239.1°C. HPLC (C18 column,

10 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 93.7%, RT = 2.10 min., <sup>1</sup>H NMR (400MHz, DMSO) δ 7.54(t, 1H, J=8.0 Hz), 7.78(d, 1H, J=7.8 Hz), 7.85(m, 3H), 8.11(d, 1H, J=8.3 Hz), 8.19(d, 1H, J=9.0 Hz), 8.53(s, 1H), 8.81(d, 1H, J=8.3 Hz), 9.14(s, 1H), 10.44(s, 1H), 11.09(s, 1H).

## EXAMPLE 131

3-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]hydrazide



Synthesis of 3-quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-

15 hydrazide: Example 131 was synthesized in accordance with the methods of Example 128 except that 3-quinolinecarboxylic acid hydrazide was used instead of 6-quinolinecarboxylic acid hydrazide.


20 MS: 362.8 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SCl); mp: 249.5-250.4°C. HPLC (C18 column,

1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.0%, RT = 2.7 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.56(s, 1H), 7.70-8.00(m, 4H), 8.10-8.20(m, 3H), 9.14(s, 1H), 10.48(s, 1H), 11.15(s, 1H).

- 113 -

## EXAMPLE 132

6-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-  
5 hydrazide: Example 132 was synthesized in accordance with the methods of  
Example 128 except that 3-fluorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 346.35 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF); mp: 250.6-251.3°C. HPLC (C18 column,  
1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 94.7%, RT = 1.8 min., <sup>1</sup>H NMR (400MHz, DMSO) δ  
10 7.40-7.70(m, 4H), 7.80(s, 1H), 8.09(d, 1H, J=7.6 Hz), 8.15(d, 1H, J=9.5 Hz), 8.51(s,  
1H), 8.76(d, 1H, J=10.5 Hz), 9.12(s, 1H), 10.40(s, 1H), 11.05(s, 1H).

## EXAMPLE 133

3-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]hydrazide




15 Synthesis of 3-quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-  
hydrazide: Example 133 was synthesized in accordance with the methods of  
Example 132 except that 3-quinolinecarboxylic acid hydrazide was used instead of  
6-quinolinecarboxylic acid hydrazide.

MS: 346.35 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF); mp: 247.8-248.4°C. HPLC (C18 column,  
20 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.7%, RT = 2.5 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
7.40-7.80 (m, 5H), 7.89(t, 1H, J=7.6 Hz), 8.00-8.20(m, 2H), 8.78(s, 1H), 9.10(s, 1H),  
10.43(s, 1H), 11.10(s, 1H).

- 114 -

## EXAMPLE 134

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide



Synthesis of 6-quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)-


5 sulfonyl]hydrazide: Example 134 was synthesized in accordance with the methods of Example 128 except that 2-(trifluoromethyl)phenylsulfonyl chloride was used instead of 3-methylbenzenesulfonyl chloride.

MS: 396.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>), mp: 236.7-237.4°C; HPLC (C18 column, 1:1MeCN/H<sub>2</sub>O + 0.1% TFA) 99.8%, RT = 1.96 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ

10 7.60-7.80(m, 3H), 7.94(d, 1H, J=6.9 Hz), 8.01(d, 1H, J=8.3 Hz), 8.10(d, 1H, J=9.1 Hz), 8.19(d, 1H, J=7.1 Hz), 8.45(S, 1H), 8.65(d, 1H, J=8.8 Hz), 9.08(S, 1H), 10.13(S, 1H), 11.06(S, 1H).

## EXAMPLE 135

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide



15

Synthesis of 3-quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)-

sulfonyl]hydrazide: Example 135 was synthesized in accordance with the methods of Example 134 except that 3-quinolinecarboxylic acid hydrazide was used instead of 6-quinolinecarboxylic acid hydrazide.

20 MS: 396.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>), mp: 203.6-204.4°C; HPLC (C18 column, 1:1


MeCN/H<sub>2</sub>O + 0.1% TFA) 93.2%, RT = 2.88 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ

7.40-8.0(m, 3H), 8.23(S, 1H), 8.67(S, 1H), 9.10(S, 1H), 10.24(S, 1H), 11.18(S, 1H).

- 115 -

## EXAMPLE 136

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide: Example 136 was synthesized in accordance with the methods of  
Example 128 except that 2-(trifluoromethoxyl)phenylsulfonyl chloride was used  
instead of 3-methylbenzenesulfonyl chloride.

MS: 412.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>4</sub>SF<sub>3</sub>), mp: 243.5-243.9°C; HPLC (C18 column,  
1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 92.1%, RT = 2.05 min., <sup>1</sup>H NMR (400 MHz, DMSO)  
δ 7.40-7.55(m, 2H), 7.60-7.80(m, 2H), δ 7.90-8.20(m, 3H), 8.43(S, 1H), 8.59(d, 1H,  
J=8.1 Hz), 9.03(S, 1H), 10.32(S, 1H), 11.00(S, 1H).

## EXAMPLE 137

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide




Synthesis of 3-quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide: Example 137 was synthesized in accordance with the methods of  
Example 136 except that 3-quinolinecarboxylic acid hydrazide was used instead of  
6-quinolinecarboxylic acid hydrazide.

MS: 412.4 (M+1 for C<sub>17</sub>H<sub>12</sub>N<sub>3</sub>O<sub>4</sub>SF<sub>3</sub>), mp: 175.1-175.9°C; HPLC (C18 column, 1:1  
MeCN/H<sub>2</sub>O + 0.1% TFA) 91.2%, RT = 2.97 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
7.40-8.10(m, 8H), 8.70(S, 1H), 9.06(S, 1H), 10.39(S, 1H), 11.08(S, 1H).

- 116 -

## EXAMPLE 138

6-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-  
5 hydrazide: Example 138 was synthesized in accordance with the methods of  
Example 128 except that 3,5-dichlorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 397.3 (M+1 for C<sub>16</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>SCl<sub>2</sub>); mp: 241.6-242.3°C. HPLC (C18 column,  
1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 91.9%, RT = 2.2 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
10 7.81(S, 3H), 7.96(S, 1H), 8.18(d, 1H, J=9.1 Hz), 8.21(d, 1H, J=9.0 Hz), 8.54(S, 1H),  
8.80(d, 1H, J=8.1 Hz), 9.13(S, 1H), 10.65(S, 1H), 11.14(S, 1H).

## EXAMPLE 139

3-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide




15 Synthesis of 3-quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-  
hydrazide: Example 139 was synthesized in accordance with the methods of  
Example 138 except that 3-quinolinecarboxylic acid hydrazide was used instead of  
6-quinolinecarboxylic acid hydrazide.

MS: 397.3 (M+1 for C<sub>16</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>SCl<sub>2</sub>); mp: 241.3-241.9°C. HPLC (C18 column,  
20 1:1 MeCN/H<sub>2</sub>O+0.1% TFA) 94.5%, RT = 4.0 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
7.75(t, 2H, J=7.1 Hz), 7.83(S, 2H), 7.96(S, 1H), 8.10-8.30(m, 2H), 8.92(S, 1H),  
9.19(S, 1H), 10.67(S, 1H), 11.23(S, 1H).

- 117 -

## EXAMPLE 140

6-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-  
sulfonyl]hydrazide: Example 140 was synthesized in accordance with the methods of  
Example 128 except that 3-chloro-4-fluorophenylsulfonyl chloride was used instead of  
3-methylbenzenesulfonyl chloride.

MS: 380.8 (M+1 for C<sub>16</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>SClF). mp: 241.9-242.6°C; HPLC (C18 column,  
1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 93.60%, RT = 2.09 min., <sup>1</sup>H NMR (400 MHz, DMSO)  
δ 7.58(t, 1H, J=9.1 Hz), 7.70-7.90(m, 2H), 8.80-8.20(m, 3H), 8.52(s, 1H), 8.78(d, 1H,  
J=8.8 Hz) 9.13(s, 1H), 10.48(s, 1H), 11.09(s, 1H).

## EXAMPLE 141

3-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide




Synthesis of 3-quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-  
sulfonyl]-hydrazide: Example 141 was synthesized in accordance with the methods of  
Example 140 except that 3-quinolinecarboxylic acid hydrazide was used instead of  
6-quinolinecarboxylic acid hydrazide.

MS: 380.8 (M+1 for C<sub>16</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>SClF). mp: 245.0-245.7°C; HPLC (C18 column,  
1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 95.69%, RT = 3.09 min., <sup>1</sup>H NMR (400 MHz, DMSO)  
δ 7.59(t, 1H, J=8.7 Hz), 7.89(t, 1H, J=5.6 Hz), 7.70-7.90(m, 2H), 8.80-8.20(m, 3H),  
8.81(d, 1H, J=8.8 Hz) 9.13(s, 1H), 10.51(s, 1H), 11.14(s, 1H).

- 118 -

## EXAMPLE 142

6-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-

5 hydrazide: Example 142 was synthesized in accordance with the methods of Example 128 except that 3-methoxyphenylsulfonyl chloride was used instead of 3-methylbenzenesulfonyl chloride.

MS: 358.4 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub>S). mp: 201.9-202.7°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 97.64%, RT = 1.73 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
10 3.73(S, 3H), 7.10-7.20(m, 2H), 7.30-7.50(m, 2H), 7.85(t, 1H, J=4.7 Hz), 8.19(d, 1H, J=8.7 Hz), 8.21(d, 1H, J=8.7 Hz), 8.59(S, 1H), 8.84(d, 1H, J=7.3 Hz), 9.17(S, 1H), 10.21(S, 1H), 11.01(S, 1H).

## EXAMPLE 143

3-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide



Synthesis of 3-quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-


15 hydrazide: Example 143 was synthesized in accordance with the methods of Example 142 except that 3-quinolinecarboxylic acid hydrazide was used instead of 6-quinolinecarboxylic acid hydrazide.

20 MS: 358.4 (M+1 for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub>S). mp: 226.9-227.2°C; HPLC (C18 column, 1:1 MeCN / H<sub>2</sub>O + 0.1 % TFA) 96.70%, RT = 2.39 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ 3.72(S, 3H), 7.10-7.20(m, 1H), 7.30-7.50(m, 2H), 7.70(t, 1H, J=7.3 Hz), 7.89(t, 1H, J=8.0 Hz), 8.00-8.20(m, 3H), 8.77(S, 1H), 9.09(S, 1H), 10.24(S, 1H), 11.03(S, 1H).

- 119 -

## EXAMPLE 144

6-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide




Synthesis of 6-quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-  
 5 hydrazide: Example 144 was synthesized in accordance with the methods of  
 Example 128 except that 3-bromophenylsulfonyl chloride was used instead of  
 3-methylbenzenesulfonyl chloride.

MS: 407.3 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SBr). mp: 216.0-216.8°C; HPLC (C18 column, 1:1  
 MeCN/H<sub>2</sub>O + 0.1% TFA) 96.88%, RT = 2.00 min., <sup>1</sup>H NMR (400 MHz, DMSO) δ  
 10 7.47(t, 2H, J=8.1), 7.70-7.90(m, 2H), 7.97(S, 1H), 8.00-8.20(m, 2H), 8.49(S, 1H),  
 8.73(d, 1H, J=8.0 Hz) 9.11(S, 1H), 10.43(S, 1H), 11.06(S, 1H).

## EXAMPLE 145

3-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]hydrazide




15 Synthesis of 3-quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-  
 hydrazide: Example 145 was synthesized in accordance with the methods of  
 Example 144 except that 3-quinolinecarboxylic acid hydrazide was used instead of  
 6-quinolinecarboxylic acid hydrazide.

MS: 407.3 (M+1 for C<sub>16</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>SBr). mp: 231.5-232.3°C; HPLC (C18 column, 1:1  
 20 MeCN/H<sub>2</sub>O + 0.1% TFA) 95.54%, RT = 2.95 min. <sup>1</sup>H NMR (400MHz, DMSO) δ  
 7.48(t, 2H, J=8.1), 7.71(t, 1H, J=8.1 Hz), 7.80-7.90(m, 2H), 7.98(S, 1H), 8.00-8.20(m,  
 2H), 8.80(S, 1H), 9.11(S, 1H), 10.47 (S, 1H), 11.13(S, 1H).


- 120 -

## EXAMPLE 146

## Benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



Step 1. The preparation of Benzofuran-2-carboxylic acid hydrazide



5 Benzofuran-2-carboxylic acid (3.85 g, 23.74 mmol) was dissolved in THF (144 mL). *n*-Methyl morpholine (5.74 mL, 52.24 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (3.39 mL, 26.12 mmol) was added, and the reaction was allowed to stir for 10 minutes. A solution of hydrazine (14.9 mL,  
10 474.8 mmol) in THF (19 mL) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1 hour. The solution was then washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was triturated with acetone to yield the desired product (2.37 g, 56.7%).

15 MS: 176.9 (M+1 for C<sub>9</sub>H<sub>8</sub>N<sub>2</sub>O<sub>2</sub>); mp: 164-167°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.23

(1:1 hexane/EtOAc). Analysis (C<sub>9</sub>H<sub>8</sub>N<sub>2</sub>O<sub>2</sub>) (calc) C: 61.36, H: 4.58, N:

15.90; (found) C: 60.96, H: 4.46, N: 15.57. IR (KBr, cm<sup>-1</sup>) 3324, 3182, 2984, 1598.

<sup>1</sup>H NMR (400 MHz, DMSO) δ 4.53 (s, 2H), 7.28 (t, 1H, J = 6.8 Hz), 7.40 (t, 1H, J = 6.8 Hz), 7.47 (s, 1H), 7.60 (d, 1H, J = 8.1 Hz), 7.72 (d, 1H, J = 7.6 Hz), and 9.99 (s,  
20 1H).

Step 2. The preparation of Benzofuran-2-carboxylic acid 2-(phenylsulfonyl)-hydrazide: Benzofuran-2-carboxylic acid hydrazide (0.4 g, 2.27 mmol) was suspended in THF (25 mL). *N,N*-diisopropylethyl amine (1.58 mL, 9.08 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.32 mL, 2.50 mmol) was  
25 added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was orange after it stirred for 1 hour, and DMF (20 mL) was added and the solution became clear. After 4 hours of stirring, the

- 121 -


solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3:1 hexane/EtOAc. The desired product was isolated (0.38 g, 53.0%).

5 MS: 317.0 (M+1 for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.23 (1:1 hexane/EtOAc); mp: 175-180°C; HRMS: 317.0598 (M+1 for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S), HPLC: (30% H<sub>2</sub>O/70% CH<sub>3</sub>CN/0.1% TFA), Ret. Time: 2.911, Purity: 93.12%. IR (KBr, cm<sup>-1</sup>) 3338, 3131, 1676, 1581, 1349, 1163. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.23-7.67 (m, 9H), 7.93 (d, 2H, J = 5.6 Hz), and 8.52 (s, 1H).


10

## EXAMPLE 147

7-Ethoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-hydrazide



Step 1. The preparation of 7-ethoxy-benzofuran-2-carboxylic acid hydrazide



15

7-Ethoxy-benzofuran-2-carboxylic acid (3.0 g, 14.5 mmol) was dissolved in THF (88 mL). N-Methyl morpholine (3.51 mL, 31.9 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (2.07 mL, 15.95 mmol) was added, and the reaction was allowed to stir for 15 minutes. Hydrazine (9.11 mL, 290 mmol) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1 hour. The solution was added EtOAc and was then washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was triturated with CH<sub>2</sub>Cl<sub>2</sub> and was then recrystallized in hot methanol (50 mL) and water to yield the desired product (2.63 g, 82.4%).

20

- 122 -

MS: 221.0 (M+1 for C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>); mp: 129-130°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.15

(1:1 hexane/EtOAc); Analysis (C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>·0.03 H<sub>2</sub>O) (calc) C: 59.85, H: 5.51, N:

12.69; (found) C: 59.51, H: 5.46, N: 12.46. IR (KBr, cm<sup>-1</sup>) 3264, 1587, 1202. <sup>1</sup>H

NMR (400 MHz, DMSO) δ 1.38 (t, 3H, J = 6.8 Hz), 4.21 (q, 2H, J = 6.8 Hz), 4.53 (s, 2H), 6.98 (dd, 1H, J = 0.7, 8.0 Hz), 7.16 (t, 1H, 8.0 Hz), 7.24 (dd, 1H, J = 1.0, 7.8 Hz), 7.44 (s, 1H), and 9.94 (s, 1H).

Step 2. The preparation of 7-ethoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-hydrazide: 7-Ethoxy-benzofuran-2-carboxylic acid hydrazide (0.5 g, 2.27 mmol) was dissolved in THF (25 mL). N,N-diisopropylethylamine (1.58 mL, 9.08 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.32 mL, 2.50 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was orange after it stirred for 1 hour. After 3 hours of stirring, the solution was added with EtOAc and washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 1:1 hexane/EtOAc. The desired product was isolated (0.05 g, 5.6%).

MS: 361.0 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.17 (1:1 hexane/EtOAc); mp:


223-224°C. HRMS: (calc) 361.0558, (found) 361.0853 (M+1 for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>S)

HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 2.596, Purity: 98.71%. IR

(KBr, cm<sup>-1</sup>) 3268, 3124, 2982, 1666, 1586, 1344, 1166, 1087, 724. <sup>1</sup>H NMR (400 MHz, DMSO) δ 1.37 (t, 3H, J = 6.8 Hz), 4.19 (q, 2H, J = 7.1 Hz), 7.01 (d, 1H, J = 8.0 Hz), 7.16-7.26 (m, 2H), 7.49-7.62 (m, 4H), 7.81 (d, 1H, J = 7.3 Hz), 10.17 (s, 1H), and 10.92 (s, 1H).


#### EXAMPLE 148

7-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



- 123 -

Step 1. The preparation of 7-Chloro-benzofuran-2-carboxylic acid hydrazide



7-Chloro-benzofuran-2-carboxylic acid (1.0 g, 5.1 mmol) was dissolved in THF (31 mL). N-Methyl morpholine (1.2 mL, 11.2 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (0.73 mL, 5.6 mmol) was added, and the reaction was allowed to stir for 10 minutes. Hydrazine (3.2 mL, 102 mmol) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1 hour. The solution was added acetate and was washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was triturated with CH<sub>2</sub>Cl<sub>2</sub>. The light brown precipitate was then recrystallized in hot methanol (25 mL) and water to yield the desired product (0.94 g, 87.8%).

MS: 211.0 (M+1 for C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Cl); mp: 179-180°C. TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.11

(1:1 hexane/EtOAc); Analysis (C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Cl·0.005H<sub>2</sub>O) (calc) C: 51.28, H: 3.36, N:

13.29; (found) C: 50.91, H: 3.44, N: 12.99. IR (KBr, cm<sup>-1</sup>) 3307, 3222, 3029, 1592, 1300, 993. <sup>1</sup>H NMR (400 MHz, DMSO) δ 4.60 (s, 2H), 7.32 (t, 1H, J = 7.9 Hz), 7.53 (d, 1H, J = 7.8 Hz), 7.59 (s, 1H), 7.73 (d, 1H, J = 7.8 Hz), and 10.10 (s, 1H).

Step 2. The preparation of 7-Chloro-benzofuran-2-carboxylic acid


2-(phenylsulfonyl)hydrazide: 7-Chloro-benzofuran-2-carboxylic acid hydrazide (0.5 g, 2.38 mmol) was suspended in THF (26 mL). N,N-diisopropylethylamine (1.66 mL, 9.52 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.33 mL, 2.62 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. The solution was yellow after it stirred for 2 hours, and DMF (10 mL) was added. After another hour of stirring, the solution was diluted with EtOAc (150 mL), and washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 3:1 hexane/EtOAc. The desired product was isolated (0.04 g, 4.6%).

- 124 -


MS: 351.0 (M+1 for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.57 (1:1 hexane/EtOAc);  
mp: 223-224°C. HRMS: (calc) 351.0206, (found) 351.0215 (M+1 for  
C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl) HPLC: Ret. Time: 9.862, Purity: 98.26%. IR (KBr, cm<sup>-1</sup>) 3291,  
3033, 1670, 1344, 1164. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24-7.64 (m, 7H), 7.94 (d,  
2H, J = 7.1 Hz), 8.62 (s, 1H), and 9.12 (s, 1H).

## EXAMPLE 149

2,3-Dihydro-benzofuran-5-carboxylic acid 2-(phenylsulfonyl)hydrazide



Step 1. The preparation of 2,3-Dihydro-benzofuran-5-carboxylic acid hydrazide



2,3-Dihydro-benzofuran-5-carboxylic acid (0.5 g, 3.05 mmol) was dissolved in  
THF (19 mL). N-Methyl morpholine (0.74 mL, 6.71 mmol) was added, and the  
reaction was cooled to 0°C. Isobutyl chloroformate (0.43 mL, 3.35 mmol) was added,  
and the reaction was allowed to stir for 10 minutes. Hydrazine (1.91 mL, 61 mmol)  
was added slowly to the reaction, and the reaction was allowed to warm to room  
temperature. The reaction was allowed to stir for 1 hour. The solution was added  
EtOAc and was washed with saturated sodium bicarbonate and brine, dried over  
Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The crude desired product was collected (0.14 g, 17.1%)  
and carried on to the next reaction.

MS: 179.1 (M+1 for C<sub>9</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.36 (1:1 hexane/EtOAc). <sup>1</sup>H  
NMR (400 MHz, DMSO) δ 3.16 (t, 2H, J = 8.8 Hz), 4.39 (s, 2H), 4.54 (t, 2H,  
J = 8.8 Hz), 6.74 (d, 1H, J = 8.3 Hz), 7.58 (d, 1H, J = 7.8 Hz), 7.67 (s, 1H), and 9.50 (s,  
1H).

Step 2. The preparation of 2,3-Dihydro-benzofuran-5-carboxylic acid

2-(phenylsulfonyl)hydrazide: 2,3-Dihydro-benzofuran-5-carboxylic acid hydrazide

- 125 -

(0.25 g, 1.40 mmol) was dissolved in THF (16 mL). N,N-diisopropylethylamine (0.98 mL, 5.6 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.20 mL, 1.54 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 2 hours of stirring, the solution was diluted with EtOAc (200 mL), and washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 1:1 hexane/EtOAc. The desired product was isolated (0.09 g, 20.0%).


MS: 319.0 (M+1 for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S); mp: 181-183°C; TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.46

(1:1 hexane/EtOAc); HRMS: (calc) 319.0752, (found) 319.0761 (M+1 for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S) HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 9.862,

Purity: 95.04%. <sup>1</sup>H NMR (400 MHz, DMSO) δ 3.13 (t, 2H, J = 8.8 Hz), 4.54 (t, 2H, J = 8.8 Hz), 6.74 (d, 1H, J = 8.3 Hz), 7.46-7.57 (m, 5H), 7.77 (d, 2H, J = 7.0 Hz), 9.85 (s, 1H), and 10.42 (s, 1H).

## EXAMPLE 150

2-Indolecarboxylic acid, 5-amino-2-phenylsulfonyl hydrazide



Example 150 was synthesized in accordance with the methods of Example 157, Step 1, except that 2-indolecarboxylic acid, 5-nitro-2-phenylsulfonyl hydrazide was used instead of 2-benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide (yield 98%).


MS: 331.1 (M+1 for C<sub>15</sub>H<sub>14</sub>N<sub>4</sub>O<sub>3</sub>S); mp: 240.2-241.1; HPLC (C18 column,

1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 95.1%, RT = 1.6 min.; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) 6.60 (d, 1H, J = 6.6 Hz), 6.64 (s, 1H), 6.87 (s, 1H), 7.05 (d, 1H, J = 8.4 Hz), 7.40-7.50 (m, 4H), 7.60 (d, 1H, J = 7.3 Hz), 7.80 (d, 2H, J = 7.0 Hz), 10.47 (s, 1H), 11.08 (s, 1H), and 11.09 (s, 1H).


- 126 -

## EXAMPLE 151

## 5-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide



Step 1. The preparation of 5-Chloro-benzofuran-2-carboxylic acid hydrazide



5-Chloro-benzofuran-2-carboxylic acid (1.0 g, 5.1 mmol) was dissolved in THF (31 mL). N-Methyl morpholine (1.2 mL, 11.2 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (0.73 mL, 5.6 mmol) was added, and the reaction was allowed to stir for 20 minutes. Hydrazine (3.2 mL, 102 mmol) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1.5 hours. The solution was added EtOAc and washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was then recrystallized in hot methanol (25 mL) and water to yield the desired product (1.07 g, 99.8%).

MS: 211.1 (M+1 for C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Cl). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.09 (1:1 hexane/EtOAc); mp: 186-187°C; Analysis (C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>2</sub>Cl) (calc) C: 51.32, H: 3.35, N: 13.30; (found) C: 51.25, H: 3.44, N: 12.90. IR (KBr, cm<sup>-1</sup>) 3321, 3186, 3020, 1666, 1599, 1442, 1315, 1183, 985, 804, 695. <sup>1</sup>H NMR (400 MHz, DMSO) δ 4.58 (s, 2H), 7.46 (d, 2H, J = 8.4 Hz), 7.67 (d, 1H, J = 8.8 Hz), 7.84 (s, 1H), and 10.08 (s, 1H).

Step 2. The preparation of 5-chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide: 5-Chloro-benzofuran-2-carboxylic acid hydrazide (0.4 g, 1.90 mmol) was suspended in THF (21 mL). N,N-diisopropylethylamine (1.32 mL, 7.6 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.27 mL, 2.09 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction

- 127 -

was allowed to warm to room temperature. The solution was orange after it stirred for 2 hours, and DMF (10 mL) was added and the solution became clear. After 15 minutes of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then

concentrated down and purified by flash chromatography eluting with

2:1 hexane/EtOAc. The desired product was isolated (0.03 g, 3.8%).

MS: 350.9 (M+1 for C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.55 (1:1 hexane/EtOAc);


mp: 181-183°C; HRMS: (calc) 351.0206, (found) 351.0206 (M+1 for

C<sub>15</sub>H<sub>11</sub>N<sub>2</sub>O<sub>4</sub>SCl) HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 8.755,


Purity: 97.13%. IR (KBr, cm<sup>-1</sup>) 3291, 3066, 1670, 1344, 1164. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 4.79 (br, 1H), 7.24-7.69 (m, 7H), 7.92 (d, 2H, J = 7.6 Hz), and 8.51 (br, 1H).

## EXAMPLE 152

Isoquinoline-1-carboxylic acid 2-(phenylsulfonyl)hydrazide



Step 1. The preparation of Isoquinoline-1-carboxylic acid hydrazide



Isoquinoline-1-carboxylic acid (2.0 g, 11.55 mmol) was dissolved in THF (70 mL). N-Methyl morpholine (2.79 mL, 25.41 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (1.65 mL, 12.7 mmol) was added, and the reaction was allowed to stir for 20 minutes. Hydrazine (7.25 mL, 231 mmol) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1.5 hours. The solution was diluted with EtOAc (200 mL) and then washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated and carried on to the next reaction.


MS: 187.9 (M+1 for C<sub>10</sub>H<sub>9</sub>N<sub>3</sub>O). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.03 (1:1 hexane/EtOAc).

- 128 -


Step 2. The preparation of isoquinoline-1-carboxylic acid 2-(phenylsulfonyl)-hydrazide: Isoquinoline-1-carboxylic acid hydrazide (2.0 g, 10.68 mmol) was dissolved in THF (119 mL). N,N-diisopropylethylamine (7.44 mL, 42.7 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (1.50 mL, 11.80 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 3 hours of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 2:1 hexane/EtOAc. The desired product was isolated (0.09 g, 2.5%).

MS: 328.1 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.48 (1:1 hexane/EtOAc); mp: 198-200°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 5.616, Purity: 98.81%. Analysis (C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S) (calc) C: 58.70, H: 4.00, N: 12.84; (found) C: 58.49, H: 3.99, N: 12.59. IR (KBr, cm<sup>-1</sup>) 3192, 1669, 1341, 1164. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.53-7.68 (m, 4H), 7.75-7.79 (m, 1H), 7.88 (d, 2H, J = 7.0 Hz), 7.94-8.00 (m, 3H), 8.48 (d, 1H, J = 5.6 Hz), 10.26 (s, 1H), and 10.82 (s, 1H).

## EXAMPLE 153

1*H*-Indole-5-carboxylic acid 2-(phenylsulfonyl)hydrazide

Step 1. The preparation of 1*H*-Indole-5-carboxylic acid hydrazide



1*H*-Indole-5-carboxylic acid (2.0 g, 12.4 mmol) was dissolved in THF (75 mL). N-Methyl morpholine (3.00 mL, 27.3 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (1.77 mL, 13.7 mmol) was added, and the reaction was allowed to stir for 20 minutes. Hydrazine (7.78 mL, 248 mmol) was

- 129 -

added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction was allowed to stir for 1.5 hours. The solution was diluted with EtOAc (200 mL) and then washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was recrystallized from hot MeOH (25 mL) and hexane (1 mL). The desired product was collected (0.52 g, 23.8%).

MS: 175.9 (M+1 for C<sub>9</sub>H<sub>9</sub>N<sub>3</sub>O); mp: 198-200°C; TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.06 (1:1 hexane/EtOAc); Analysis (C<sub>9</sub>H<sub>9</sub>N<sub>3</sub>O) (calc) C: 61.70, H: 5.18, N: 23.99; (found) C: 61.90, H: 5.29, N: 24.09. IR (KBr, cm<sup>-1</sup>) 3271, 1585, 1521. <sup>1</sup>H NMR (400 MHz, DMSO) δ 4.37 (s, 2H), 6.47 (s, 1H), 7.35-7.38 (m, 2H), 7.55 (d, 1H, J = 8.5 Hz), 8.04 (s, 1H), 9.53 (s, 1H), and 11.27 (s, 1H).


Step 2. The preparation of 1*H*-Indole-5-carboxylic acid 2-(phenylsulfonyl)-hydrazide: 1*H*-Indole-5-carboxylic acid hydrazide (0.5 g, 2.85 mmol) was suspended in THF (32 mL). N,N-diisopropylethylamine (1.99 mL, 11.40 mmol) was added and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.36 mL, 2.85 mmol) was added followed by DMAP (5 mg 0.04 mmol). The reaction was allowed to warm to room temperature, and DMF (3 mL) was added. After 3 hours of stirring, the solution was diluted with EtOAc (200 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 2:1 hexane/EtOAc. The desired product was isolated (0.41 g, 23.8%).

MS: 316.0 (M+1 for C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.38 (1:1 hexane/EtOAc); mp: 209-211°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 4.102, Purity: 96.75%. Analysis (C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S) (calc) C: 57.13, H: 4.16, N: 13.32; (found) C: 57.28, H: 4.08, N: 13.06. IR (KBr, cm<sup>-1</sup>) 3293, 3168, 1649, 1312, 1168. <sup>1</sup>H NMR (400 MHz, DMSO) δ 6.48 (s, 1H), 7.33-7.41 (m, 3H), 7.47 (t, 2H, J = 7.3 Hz), 7.57 (t, 1H, J = 7.4 Hz), 7.80 (d, 2H, J = 7.3 Hz), 7.94 (s, 1H), 9.84 (s, 1H), 10.45 (s, 1H), and 11.33 (s, 1H).

- 130 -

## EXAMPLE 154

## Quinoline-3-carboxylic acid 2-(phenylsulfonyl)hydrazide



Quinoline-3-carboxylic acid hydrazide (0.34 g, 1.82 mmol) was dissolved in THF (20 mL). N,N-diisopropylethylamine (1.27 mL, 7.28 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.23 mL, 1.82 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 2 hours of stirring, the red solution was diluted with EtOAc (100 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 2:1 hexane/EtOAc. The desired product was isolated (0.13 g, 21.8%).


MS: 328.1 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.21 (1:1 hexane/EtOAc); mp: 249-250°C. HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 2.701, Purity:

97.64%. Analysis (C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S·0.08H<sub>2</sub>O) (calc) C: 58.45, H: 4.03, N:

12.78; (found) C: 58.52, H: 4.25, N: 12.38. IR (KBr, cm<sup>-1</sup>) 3282, 3064, 2823, 1691, 1294, 1164. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.51 (t, 2H, J = 7.8 Hz), 7.58-7.68 (m, 2H), 7.83-7.86 (m, 3H), 8.02-8.08 (m, 2H), 8.68 (s, 1H), 9.02 (s, 1H), 10.18 (s, 1H), and 11.00 (s, 1H).


## EXAMPLE 155

## Quinoline-4-carboxylic acid 2-(phenylsulfonyl)hydrazide



- 131 -

Step 1. The preparation of quinoline-4-carboxylic acid hydrazide



Quinoline-4-carboxylic acid (0.75 g, 4.33 mmol) was dissolved in THF (18 mL). N-Methyl morpholine (1.05 mL, 9.52 mmol) was added, and the reaction was cooled to 0°C. Isobutyl chloroformate (0.62 mL, 4.76 mmol) was added, and the reaction was allowed to stir for 20 minutes. Hydrazine (2.72 mL, 86.6 mmol) was added slowly to the reaction, and the reaction was allowed to warm to room temperature. The reaction stirred for 1.5 hours. The solution was diluted with EtOAc (100 mL) and then washed with saturated sodium bicarbonate and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The precipitate was carried on to the next reaction.

MS: 187.9 (M+1 for C<sub>10</sub>H<sub>9</sub>N<sub>3</sub>O). <sup>1</sup>H NMR (400 MHz, DMSO) δ 4.64 (s, 2H), 7.45-7.48 (m, 1H), 7.60-7.65 (m, 1H), 7.75-7.79 (m, 1H), 8.01-8.15 (m, 2H), 8.90-8.93 (m, 1H), and 9.89 (s, 1H).

Step 2. The preparation of quinoline-4-carboxylic acid 2-(phenylsulfonyl)-hydrazide:

Quinoline-4-carboxylic acid hydrazide (0.6 g, 3.25 mmol) was dissolved in THF (36 mL). N,N-diisopropylethylamine (2.26 mL, 13.0 mmol) was added, and the reaction was cooled to 0°C. Phenylsulfonyl chloride (0.41 mL, 3.25 mmol) was added followed by DMAP (5 mg, 0.04 mmol). The reaction was allowed to warm to room temperature. After 2.5 hours of stirring, the red solution was diluted with EtOAc (150 mL). It was then washed with saturated sodium bicarbonate and brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The reaction was then concentrated down and purified by flash chromatography eluting with 1:1 hexane/EtOAc. The desired product was isolated (0.05 g, 18.8%).

MS: 328.0 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S). TLC: SiO<sub>2</sub>, R<sub>f</sub> = 0.10 (1:1 hexane/EtOAc); mp: 153-156°C; HRMS: (calc) 328.0756, (found) 328.0753 (M+1 for C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>S) HPLC: (50% H<sub>2</sub>O/50% CH<sub>3</sub>CN/0.1% TFA) Ret. Time: 2.284, Purity: 100.0%. IR (KBr, cm<sup>-1</sup>) 3199, 2963, 1745, 1665, 1168. <sup>1</sup>H NMR (400 MHz, DMSO) δ 7.43 (d,

- 132 -


1H, J = 4.2 Hz), 7.49-7.80 (m, 8H), 7.91 (d, 1H, J = 7.1 Hz), 8.01-8.07 (m, 1H), 8.94 (d, 1H, J = 3.9 Hz), 10.36 (s, 1H), 10.93 (s, 1H).

Additional compounds that can be prepared according to the scheme and examples set forth above:

- 5           1. Dibenzofuran-2-carboxylic acid 2-(4-fluoro-3-methylphenylsulfonyl)-hydrazide
2. 7-Methoxy-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide
3. 7-Chloro-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide
4. 5-Methoxy-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide
- 10          5. 5-Chloro-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide
6. 5-Chloro-1*H*-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-phenylsulfonyl]hydrazide
7. 5-Methyl-1*H*-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide
8. 5-Methyl-1*H*-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-phenylsulfonyl]hydrazide
- 15

## EXAMPLE 156

2-Indolecarboxylic acid, [(5-N, N-dimethylamino)-, 2-phenylsulfonyl] hydrazide hydrochloride salt



- 20           Synthesis of 2-indolecarboxylic acid, [(5-N,N-dimethylamino), 2-phenylsulfonyl] hydrazide hydrochloride salt: Example 156 was synthesized in accordance with the methods of example 167 except that 2-indolecarboxylic acid, 5-amino-, 2-phenylsulfonyl hydrazide was used instead of dibenzofurancarboxylic acid, 2-amino-, 2-(phenylsulfonyl)hydrazide (yield 94%).
- 25           MS: 359.1 (M+1 for C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O<sub>3</sub>S); mp: 251.6-252.7°C; HPLC (C18 column, 1:1 MeCN / H<sub>2</sub>O = 0.1%TFA) 82.4%, RT=2.2 min; <sup>1</sup>HNMR(DMSO-D<sub>6</sub>) δ 2.46(S,


- 133 -

6H), 7.51(t, 2H, J=7.8Hz), 7.62(t, 1H, J=7.2Hz), 7.68(s, 1H), 7.70-7.90(m, 4H), 8.09(s, 1H), 10.23(s, 1H), 11.03(s, 1H)

## EXAMPLE 157

2-Benzofurancarboxylic acid, 5-(phenylsulfonylamino)-, 2-(phenylsulfonyl) hydrazide


5 Step 1. 2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide



This was prepared in accordance with the following method.

2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide (0.82 g) was dissolved in 50 mL of THF:EtOH (1:1), treated with wet Raney Ni (0.6 g) and hydrogenated at 50 Psi for 15 hours. The solvent was removed to give 0.73 g (97%) of the desired product as a white solid.

Step 2. 2-Benzofurancarboxylic acid, 5-(phenylsulfonylamino)-, 2-(phenylsulfonyl)hydrazide




15 2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide (0.117 g, 0.35 mmol) was dissolved in 7 mL of dry pyridine, then benzenesulfonyl chloride (0.03 mL, 0.39 mmol) was added, the reaction was stirred at RT overnight. The solvent was removed and recrystallized from ethyl acetate and hexane to give 0.155 g (94%) of the desired product as a white solid.

20 MS: 472.0 (M+1 for C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>S<sub>2</sub>); mp: 261.5-261.8°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O = 0.1%TFA) 96.5%, RT = 6.0 min.; <sup>1</sup>HNMR(DMSO-D<sub>6</sub>) δ 7.10 (d, 1H, J = 6.8Hz), 7.41(s, 1H), 7.40-7.60 (m, 8H), 7.67 (d, 2H, J = 7.5 Hz), 7.79 (d, 2H, J=7.3Hz), 10.15(s, 1H), 10.25(s, 1H), 10.89(s, 1H).

- 134 -

## EXAMPLE 158

2-Benzofurancarboxylic acid, 5-(methylsulfonylamino)-, 2-(phenylsulfonyl) hydrazide



Synthesis of 2-benzofurancarboxylic acid, 5-(methylsulfonylamino)-,


5 2-(phenylsulfonyl)hydrazide: Example 158 was synthesized in accordance with the methods of Example 157 except that methylsulfonyl chloride was used instead of benzenesulfonyl chloride (yield 92%).

MS: 410.0 (M+1 for C<sub>16</sub>H<sub>15</sub>N<sub>3</sub>O<sub>6</sub>S<sub>2</sub>); mp: 255.4-256.1°C; HPLC (C18 column,

10 1:1 MeCN / H<sub>2</sub>O = 0.1%TFA) 90.0%, RT = 3.3 min.; <sup>1</sup>HNMR(DMSO-D<sub>6</sub>) δ 2.92 (s, 3H), 7.26 (d, 1H, J = 6.8Hz), 7.50-7.60 (m, 6H), 7.81 (d, 1H, J = 7.6 Hz), 9.60 (s, 1H), 9.70 (s, 1H), 10.18 (s, 1H), 10.93 (s, 1H).

## EXAMPLE 159

2-Dibenzofurancarboxylic acid, 2-(4-nitrophenyl)sulfonyl hydrazide



15 Synthesis of 2-dibenzofurancarboxylic acid, 2-(4-nitrophenyl)sulfonyl hydrazide: Example 159 was synthesized in accordance with the methods of Example 92 except that 4-nitrobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride (yield 93%).


MS: 412.0 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>3</sub>O<sub>6</sub>S): mp: 239.3-239.9°C; HPLC (C18 column,

20 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 98.5%, RT = 14.52 min; <sup>1</sup>HNMR(DMSO-D<sub>6</sub>) δ 7.40 (t, 1H, J = 8.30 Hz), 7.53 (t, 1H, J = 7.33 Hz), 7.73 (m, 2H), 7.85 (dd, 1H, J = 8.55 and 1.71 Hz), 8.11-8.17 (m, 3H) 8.36 (d, 2H, J = 5.86 Hz), 8.50 (s, 1H), 10.55 (s, 1H), 10.89 (s, 1H).

- 135 -

## EXAMPLE 160

2-Benzofurancarboxylic acid, 5-(4-methoxyphenylsulfonylamino)-,  
2-(phenylsulfonyl)hydrazide




Synthesis of 2-benzofurancarboxylic acid, 5-(4-methoxyphenylsulfonyl-aminoamino)-, 2-(phenylsulfonyl)hydrazide: Example 160 was synthesized in accordance with the methods of Example 157 except that 4-methoxybenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (yield 93%).

MS: 502.0 (M+1 for C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>7</sub>S<sub>2</sub>); mp: 213-214°C; HPLC (C18 column,

1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 96.0%, RT = 4.8 min., <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 3.72 (s, 3H), 6.96 (d, 2H, J = 8.8 Hz), 7.14 (d, 1H, J = 7.1 Hz), 7.40-7.7 (m, 4H), 7.78 (d, 2H, J = 7.5 Hz), 7.98 (t, 2H, J = 6.8 Hz), 8.48 (t, 1H, J = 7.8 Hz), 8.88 (d, 1H, J = 5.1 Hz), 10.18 (s, 1H), 10.21 (s, 1H), 10.92 (s, 1H).

## EXAMPLE 161

2-Benzofurancarboxylic acid, 5-(4-methylphenylsulfonylamino)-, 2-(phenylsulfonyl)hydrazide



Synthesis of 2-benzofurancarboxylic acid, 5-(4-methylphenylsulfonylamino)-, 2-(phenylsulfonyl) hydrazide: Example 161 was synthesized in accordance with the methods of Example 157 except that 4-methylbenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (yield 91%).

MS: 486.0 (M+1 for C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>6</sub>S<sub>2</sub>); mp: 181-183°C; HPLC (C18 column,


1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 93.0%, RT = 5.5 min.; <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 2.26 (s, 3H), 7.07 (d, 1H, J = 7.8 Hz), 7.25 (d, 2H, J = 8.1 Hz), 7.40-7.60 (m, 4H), 7.80 (d, 2H,

- 136 -

J = 7.6 Hz), 7.99 (t, 2H, J = 6.8 Hz), 8.48 (t, 1H, J = 7.8 Hz), 8.88 (d, 1H, J = 5.1 Hz), 10.18 (s, 1H), 10.27 (s, 1H), 10.91 (s, 1H).

## EXAMPLE 162

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl] hydrazide




Synthesis of 2-dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)

sulfonyl]hydrazide: Example 162 was synthesized in accordance with the methods of Example 92 except that 2-nitrobenzenesulfonyl chloride was used instead of 2-fluorobenzenesulfonyl chloride (yield 93%).

MS: 412.0 (M+1 for C<sub>19</sub>H<sub>13</sub>N<sub>3</sub>O<sub>6</sub>S): mp: 223.4-224.1°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 93.2%, RT = 9.8 min; <sup>1</sup>HNMR (CDCl<sub>3</sub>) δ 7.42 (t, 1H, J = 8.30 Hz), 7.48 (t, 1H, J = 7.33 Hz), 7.65-7.95 (m, 6H), 8.10 (d, 1H, J = 5.86 Hz), 8.19 (d, 1H, J = 1.47 Hz), 10.38 (s, 1H), 10.94 (s, 1H).

## EXAMPLE 163

2-Benzofurancarboxylic acid, 5-(4-pentylphenylsulfonylamino)-, 2-(phenylsulfonyl)hydrazide



Synthesis of 2-benzofurancarboxylic acid, 5-(4-pentylphenylsulfonyl amino)-, 2-(phenylsulfonyl)hydrazide: Example 163 was synthesized in accordance with the methods of Example 157 except that 4-pentylbenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (yield 97%).


MS: 541.0 (M+1 for C<sub>26</sub>H<sub>27</sub>N<sub>3</sub>O<sub>6</sub>S<sub>2</sub>); mp: 169.0-169.8°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 93.2%, RT = 16.9 min.; <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 0.78 (t, 3H, J = 6.9 Hz), 1.18 (m, 4H), 1.42 (m, 2H), 2.50 (m, 2H), 7.13 (d, 1H, J = 8.3 Hz),

- 137 -

7.27 (d, 2H, J = 8.1 Hz), 7.40-7.60 (m, 4H), 7.80 (d, 2H, J = 7.6 Hz), 8.01 (t, 2H, J = 6.6 Hz), 8.52 (t, 1H, J = 7.8 Hz), 8.89 (d, 1H, J = 5.1 Hz), 10.17 (s, 1H), 10.27 (s, 1H), 10.91 (s, 1H).

## EXAMPLE 164

5 2-Benzofurancarboxylic acid, 5-(N-isopropylamino)-, 2-(phenylsulfonyl) hydrazide



2-Benzofurancarboxylic acid, 5-(N-isopropylamino)-, 2-(phenylsulfonyl)

hydrazide was prepared in accordance with the following method:


10 2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide (0.47 g) was dissolved in 20 mL of EtOH, treated with 20% Pd/C (50 mg), 3 mL of acetone and hydrogenated at 50-38 Psi for 15 hours. The solvent was removed to give 0.49 g (93%) of the desired product as a white solid.

MS: 374.1 (M+1 for C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 94.7-95.5°C; HPLC (C18 column,

15 1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 99.9%, RT = 1.7 min.; <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 1.09 (d, 6H, J = 6.4 Hz), 3.45 (m, 1H), 6.65 (s, 1H), 6.77 (d, 1H, J = 6.8 Hz), 7.29 (d, 1H, J = 8.7 Hz), 7.33 (s, 1H), 7.53 (t, 2H, J = 7.6 Hz), 7.61 (t, 1H, J = 7.4 Hz), 7.81 (d, 2H, J = 7.6 Hz), 10.08 (s, 1H), 10.09 (s, 1H), 10.75 (s, 1H).

## EXAMPLE 165

20 2-Benzofurancarboxylic acid, 5-(4-propylphenylsulfonylamino)-, 2-(phenylsulfonyl) hydrazide




25 Synthesis of 2-benzofurancarboxylic acid, 5-(4-propylphenylsulfonylamino)-, 2-(phenylsulfonyl) hydrazide: Example 165 was synthesized in accordance with the methods of Example 157 except that 4-propylbenzenesulfonyl chloride was used instead of benzenesulfonyl chloride (yield 97%).

- 138 -

MS: 514.1 (M+1 for C<sub>24</sub>H<sub>23</sub>N<sub>3</sub>O<sub>6</sub>S<sub>2</sub>); mp: 160.4-161.1°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 93.5%, RT = 9.3 min.; <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 0.80 (m, 3H), 1.56 (m, 2H), 2.50 (m, 2H), 7.12 (m, 1H), 7.27 (d, 2H, J = 8.1 Hz), 7.40-7.60 (m, 4H), 7.80 (d, 2H, J = 7.6 Hz), 8.01 (t, 2H, J = 6.6 Hz), 8.52 (t, 1H, J = 7.8 Hz), 8.89 (d, 1H, J = 5.1 Hz), 10.17 (s, 1H), 10.25 (s, 1H), 10.90 (s, 1H).

## EXAMPLE 166

2-Dibenzofurancarboxylic acid, 2-[(2-aminophenyl)sulfonyl]hydrazide



Synthesis of 2-dibenzofurancarboxylic acid, 2-[(2-aminophenyl)sulfonyl]

hydrazide: Example 166 was synthesized in accordance with the methods of Example 157, Step 1, except that 2-dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]hydrazide was used instead of 2-benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)hydrazide (yield 94%).

MS: 382.0 (M+1 for C<sub>19</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 217.9-218.4°C; HPLC (C18 column,


1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 98.9%, RT = 6.7 min.; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ 6.15 (s, 2H), 6.50 (t, 1H, J = 8.20 Hz), 6.76 (t, 1H, J = 9.0 Hz), 7.21 (t, 1H, J = 8.30 Hz), 7.40-7.60 (m, 3H), 7.72 (t, 1H, J = 8.30 Hz), 7.85 (t, 1H, J = 7.1 Hz), 8.10 (d, 1H, J = 7.4 Hz), 8.52 (s, 1H), 9.75 (s, 1H), 10.68 (s, 1H).

## EXAMPLE 167

2-Dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl)sulfonyl] hydrazide hydrochloride salt

Step 1: 2-Dibenzofurancarboxylic acid, 2-[(N,N-dimethyl-2-aminophenyl) sulfonyl] hydrazide


- 139 -



Synthesis of 2-dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl)-sulfonyl]hydrazide: This was prepared in accordance with the following method:

2-Benzofurancarboxylic acid, 2-amino-, 2-(phenylsulfonyl)hydrazide (0.1 g) was dissolved in 4 mL of EtOH, treated with 20% Pd/C (20 mg), 1 mL of formaldehyde and hydrogenated at 50-38 Psi for 16 hours. The solvent was removed to give 0.096 g (89%) of the desired product as a oily liquid.

Step 2: 2-Dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl)sulfonyl]hydrazide hydrochloride salt.



This was prepared in accordance with the following method:

2-Dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl)sulfonyl] hydrazide was dissolved in 5 mL of MeOH, then 2 mL of 1 M HCl (ether) was added slowly and the mixture was stirred at RT for 10 minutes, the solvent was removed to yield a white solid as the desired product.


MS: 410.1 (M+1 for C<sub>21</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 103.2-103.9; HPLC (C18 column,

1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 81.2%, RT = 8.1 min.; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ 2.97 (s, 6H), 6.68 (d, 1H, J = 8.8 Hz), 7.40 (t, 1H, J = 6.42 Hz), 7.50-7.80 (m, 6H), 7.84 (d, 1H, J = 8.30 Hz), 8.17 (d, 1H, J = 6.1 Hz), 8.49 (s, 1H), 10.41 (s, 1H), 10.68 (s, 1H).

- 140 -

## EXAMPLE 168

2-Benzofurancarboxylic acid, 5-(N,N-dimethylamino)-, 2-(phenylsulfonyl) hydrazide hydrochloride salt




2-Benzofurancarboxylic acid, 5-(N,N-dimethylamino)-, 2-(phenylsulfonyl) hydrazide hydrochloride salt: Example 168 was synthesized in accordance with the methods of Example 167 except that 2-benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)hydrazide was used instead of 2-dibenzofurancarboxylic acid, 2-[(2-N,N-dimethylaminophenyl) sulfonyl] hydrazide (yield 89%).


MS: 360.1 (M+1 for C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>4</sub>S); mp: 218.3-218.9°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O = 0.1% TFA) 99.7%, RT = 1.6 min.; <sup>1</sup>HNMR (DMSO-D<sub>6</sub>) δ 2.46 (s, 6H), 7.51 (t, 2H, J = 7.8 Hz), 7.62 (t, 1H, J = 7.2 Hz), 7.68 (s, 1H), 7.70-7.90 (m, 4H), 8.09 (s, 1H), 10.23 (s, 1H), 11.03 (s, 1H).

## EXAMPLE 169

2-Indolecarboxylic acid, 5-chloro-2-[(2-(trifluoromethyl)phenyl)sulfonyl]hydrazide



Step 1. The preparation of 5-chloroindole-2-carboxylic acid hydrazide



This was synthesized in accordance with the methods of Example 152 Step 1 except that 5-chloroindole-2-carboxylic acid was used instead of benzofuran-2-carboxylic acid (yield 94%).

Step 2. Synthesis of 2-Indolecarboxylic acid, 5-chloro-2-[(2-(trifluoromethyl)phenyl)sulfonyl]hydrazide was run in accordance with Example 51 except that


- 141 -

5-chloroindole-2-carboxylic acid hydrazide was used instead of 5-chloro-2-benzofurancarboxylic acid hydrazide. (86% yield).


MS: 418.0 (M+1 for C<sub>16</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>SF<sub>3</sub>Cl): mp: 66.8-67.9°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 91.2%, RT = 1.49 min; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>), 7.16 (s, 1H), 7.60-7.90 (m, 3H), 8.03 (t, 2H, J = 6.7 Hz), 8.56 (t, 1H, J = 7.8 Hz), 8.90 (d, 1H, J = 5.2 Hz), 10.08 (s, 1H), 10.89 (s, 1H), 11.83 (s, 1H).

## EXAMPLE 170

2-Indolecarboxylic acid, 5-nitro-2-phenylsulfonyl hydrazide



Step 1. The preparation of 5-nitroindole-2-carboxylic acid hydrazide



This was synthesized in accordance with the methods of Example 152, Step 1 except that 5-nitroindole-2-carboxylic acid was used instead of benzofuran-2-carboxylic acid (yield 92%).


Step 2. Synthesis of 2-indolecarboxylic acid, 5-nitro-2-phenylsulfonyl hydrazide was run in accordance with Example 51 except that 5-nitroindole-2-carboxylic acid hydrazide was used instead of 5-chloro-2-benzofurancarboxylic acid hydrazide and phenylsulfonyl chloride was used instead of 2-trifluoromethyl phenylsulfonyl chloride (86% yield).

MS: 361.1 (M+1 for C<sub>15</sub>H<sub>12</sub>N<sub>4</sub>O<sub>5</sub>Si): mp: 268.2-269.1°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1 %TFA) 89.2%, RT = 4.6 min; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>), 7.41 (s, 1H), 7.50-7.70 (m, 4H), 7.95 (d, 2H, J = 7.1 Hz), 8.12 (dd, 1H, J = 2.6 and 6.8 Hz), 8.09 (s, 1H), 10.18 (s, 1H), 10.96 (s, 1H), and 12.29 (s, 1H).


- 142 -

## EXAMPLE 171

2-Indolecarboxylic acid, 5-chloro-2-phenylsulfonyl hydrazide



Step 1. The preparation of 5-chloroindole-2-carboxylic acid hydrazide



5

This was synthesized in accordance with the methods of Example 152, Step 1 except that 5-chloroindole-2-carboxylic acid was used instead of benzofuran-2-carboxylic acid (yield 90%).

Step 2. Synthesis of 2-Indolecarboxylic acid, 5-chloro-2-phenylsulfonyl hydrazide was run in accordance with Example 51 except that 5-chloroindole-2-carboxylic acid hydrazide was used instead of 5-chloro-2-benzofurancarboxylic acid hydrazide and phenylsulfonyl chloride was used instead of 2-trifluoromethyl-phenylsulfonyl chloride (89% yield).

MS: 350.1 (M+1 for C<sub>15</sub>H<sub>12</sub>ClN<sub>3</sub>O<sub>3</sub>S): mp: 79.1-79.9°C; HPLC (C18 column, 1:1 MeCN/H<sub>2</sub>O + 0.1% TFA) 93.8%, RT = 6.1 min; <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) 7.16 (s, 1H), 7.40-7.60 (m, 3H), 7.80 (d, 1H, J = 7.6 Hz), 8.01 (t, 2H, J = 5.1 Hz), 10.18 (s, 1H), 8.90 (d, 1H, J = 5.2 Hz), 10.79 (s, 1H), 10.14 (s, 1H) and 11.82 (s, 1H).

## EXAMPLE 172

*In vitro* enzyme assay – human Branched Chain Amino Acid Aminotransferase cytosolic form (hBCATc) and mitochondrial form (hBCATm) as well as rat BCAT cytosolic form were assayed at 37°C in 25 mM phosphate buffer pH 7.8. In addition, 2 mM DTT and 12.5 mM EDTA were added to the assay mixture. A coupling enzyme assay was used to monitor the production of glutamate. The formation of NADH from NAD<sup>+</sup> at 340 nM was followed on a 96-well plate Molecular Devices plate reader. The following components were used in this coupled assay: 4 mM ADP, 1 mM

- 143 -

NAD<sup>+</sup>, 750  $\mu$ M L-Leucine, 500  $\mu$ M  $\alpha$ -ketoglutarate, 10  $\mu$ M pyridoxal phosphate, 1 unit glutamate dehydrogenase, and 1.25  $\mu$ g of the appropriate BCAT enzyme. Inhibitions by compounds were assayed by adding various concentrations to this coupled assay procedure as a DMSO stock up to a 5% v/v DMSO/buffer ratio.

5           *Enzyme Preparation-* humanBCATc and ratBCATc were expressed in BL21(DE3) cells. A growing culture of cells was induced with 1 mM IPTG at room temperature overnight. These cells were harvested by centrifugation at 10,000 g for 20 minutes. The supernatant was discarded, and the cells were stored at -80°C until needed. The protein was purified by the following method. The cell paste was  
10           resuspended in extraction buffer (0.1 M phosphate, pH 8.0, with 0.01 M Tris-HCL and 5 mM TCEP and then lysed on a French Press at 1000 psi. The lysate was centrifuged at 10,000 g for 15 minutes, and the pellet was discarded. The supernatant was loaded onto a Hitrap Chelating column previously charged with 0.1 M NiSO<sub>4</sub> and washed with 3 to 4 column volumes of extraction buffer. The column was then washed with  
15           0.01 Tris, pH 7.5, 10% glycerol, 150 mM NaCl, 5 mM TCEP for two column volumes. The column was then further washed with two column volumes of 0.1 M phosphate, pH 6.0, 0.01 M Tris, 10% glycerol and 750 mM NaCl. Finally, the column was washed with two column volumes of the same buffer plus 50 mM imidazole. The BCAT enzyme was then eluted with the same buffer but with 350 mM imidazole. The eluant  
20           was then dialyzed overnight against 10 mM phosphate buffer pH 8.0, 10 % glycerol, and 5 mM TCEP. The dialyzed protein was then loaded onto a Q-sepharose column and eluted using a salt gradient. Active fractions were collected and further purified on a Superose-12 column to yield pure BCAT protein.

- 144 -

TABLE 1

In-Vitro Data

Example	RrBCATc ( $\mu$ M)	RhBCATc ( $\mu$ M)
1	0.18	2.9, 1.4
2	0.13	2.6
3	2.5	17.8
4	1.22	12.9
5	0.86, 0.46, 0.53	5.76, 3.0
6	0.66	12.9
7	0.59	11.5
8	0.3	3.0
9	2.18, 3.3	17.9, 29
10	2.9	65.6
11	--	--
12	--	90.5
13	0.39	15.7
14	0.37, 0.14	2.2, 1.4
15	0.65, 0.22	3.6, 3.3
16	0.39, 0.20	3.0, 3.1
17	1.1, 0.54	11.4
18	0.53, 0.4	7.9
19	0.22	6.7
20	0.39	15.7
21	--	>100
22	4.1	24.6
23	5.4	17.80
24	1.20	24
25	1.8	21.9
26	15.20, 6.90	138.0
27	45.2	171
28	--	>100
29	0.29	10.1

- 145 -

TABLE 1 (cont'd)

## In-Vitro Data

Example	RrBCATc ( $\mu\text{M}$ )	RhBCATc ( $\mu\text{M}$ )
30	5.7	25
31	0.81	2.60, 4.30
32	>100	>100
33	3.3	12.2
34	0.33	5.1
35	0.31	3.57
36	1.1	17.5, 18.20
37	3.5	41.3
38	0.77, 0.61	29.9
39	0.21	2.29
40	1.00	4.86, 8.80
41	14.5	28.8
42	1.3	5.3, 10.40
43	0.47	1.8
44	--	3.44
45	0.21	2.29
46	--	1.08, 0.7
47	0.15, 0.2	0.84, 1.47
48	0.91	4.21, 7.60
49	3.1, 4.0	1.75, 3.60
50	1.3	5.18, 9.10
51	0.072, 0.14	0.79, 0.82
52	0.3	2.19
53		
54	--	3.76
55	--	3.53, 3.3
56	<0.50	7.26
57	0.14	1.14
58	0.22	3.57

- 146 -

TABLE 1 (cont'd)

## In-Vitro Data

Example	RrBCATc ( $\mu\text{M}$ )	RhBCATc ( $\mu\text{M}$ )
59	0.12	3.42
60	0.11	21.10, 12.8
61	0.1	1.3
62	0.33	4.77
63	5.29	66.3
64	0.19	3.31
65	0.1	3.23
66	0.15	1.38
67	2.4	17.3
68	0.31	2.78
69	0.05	3.86
70	0.43	11.62
71	0.88	6.57
72	1.86	>10, 5.35
73	1.89	14.42
74	3.71	64.5
75	0.88	>10, 5.4
76	1.51	11.24
77	0.35	6.15, 3.0
78		3.70
79	0.22	16.9, 1.81
80	0.25	2.62
81	0.31	1.74
82	0.51	3.28
83	0.87	9.66
84	0.97	6.13
85	1.39	21.18
86	0.24	15.2
87	1.16	41.07

- 147 -

TABLE 1 (cont'd)

## In-Vitro Data

Example	RrBCATc ( $\mu\text{M}$ )	RhBCATc ( $\mu\text{M}$ )
88	4	40
89	>100	>100
90	>100	>100
91	0.11	19.6
92	1.1	12.1
93	5.9	63.5
94	0.72	5.5
95	0.3	2.9
96	36.3	51.6
97	36.4	67.5
98	0.65	2.8
99	35.1	46.3
100	4.0	28.1
101	22	48.1
102	0.37	6.0
103	0.85	12
104	1.4	23
105	3.3	54
106	5.6	42
107	7.5	41
108	5.6	57
109	0.46	9.8
110	0.72	11.1
111	0.47	0.47, 2.8
112	6.2	14.6
113	0.53	4.1
114	26.5	59.7
115	5.2	59.9
116	1.2	8.2

- 148 -

TABLE 1 (cont'd)

## In-Vitro Data

Example	RrBCATc ( $\mu$ M)	RhBCATc ( $\mu$ M)
117	0.82	5.7
118	1.8	25.0
119	2.2	12.7
120	2.0	18.1
121	11.6	55.0
122	56.7	58.0
123	2.2	11.6
124	100	53.1
125	2.4	18.4
126	12.5	36.2
127		16.8
128		12
129		13
130		22
131		61
132		40
133		>100
134		17
135	2.8	13
136		40
137	14	41
138		>100
139		41
140		>100
141		>100
142		46
143		80
144		29
145		80

- 149 -

TABLE 1 (cont'd)


## In-Vitro Data

Example	RrBCATc ( $\mu$ M)	RhBCATc ( $\mu$ M)
146	2.5	17.8
147	--	12.9
148	0.59	11.5
149	3.3	17.9
150		
151	0.3	3
152	--	>100
153	1.9	15.3
154	3.6	24.2
155	--	>100
156		
157		8.7
158		26
159	4	28
160		
161		
162		
163		
164		
165		
166		
167		
168		
169		
170		
171		

- 150 -

CLAIMS


1. A compound of Formula I



wherein:

- 5  $R_3$  is H, or F, Br, alkyl, carboxy, alkoxy, substituted alkoxy,  
 $R_1$ ,  $R_2$ ,  $R_4$ , and  $R_5$  are independently, H, halogen, alkyl, substituted alkyl,  
 alkoxy, substituted alkoxy, cyano, nitro, amino, alkylamine, and  
 thioalkyl, Ar is bicyclic heteroaryl, tricyclic heteroaryl, substituted  
 bicyclic heteroaryl, except (6-methoxybenzofuran)-2-nyl, 3-quinolinyll;  
 10 or a pharmaceutically acceptable salt, ester, prodrug, or amide thereof.

2. A compound according to Claim 1 wherein Ar is selected from the group consisting of:



15

- 151 -

wherein R<sub>6</sub> is alkoxy, substituted alkoxy, halogen, aryloxy, substituted aryloxy, alkylamino, substituted alkylamino, arylalkylamino, substituted arylalkylamino, amide or substituted amide.

3. A compound according to Claim 1 wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub> are hydrogen.

4. A compound according to Claim 1 wherein Ar is a tricyclic heteroaryl.

5. A compound according to Claim 1 wherein Ar is a bicyclic heteroaryl.

6. A compound according Claim 1 wherein the compound is:

Dibenzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2,3,4,9-Tetrahydro-1H-carbazole-6-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

Benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Ethoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

Benzo[1,3] dioxole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1-Dibenzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

- 152 -

2-Dibenzofurancarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(3,5-dichlorophenylsulfonyl)-hydrazide;

5 Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-8-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(4-fluorophenyl)]-sulfonyl hydrazide;

10 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-fluorophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-fluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-chloro-4-fluorophenyl)sulfonyl]hydrazide;

15 2-Dibenzofurancarboxylic acid, 2-[(3-cyanophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-cyanophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2,6-dichlorophenyl)sulfonyl]-hydrazide;

20 2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(2-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide;

25 2-Dibenzofurancarboxylic acid, 2-[[3,5-bis(trifluoromethyl)-phenyl]-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)-hydrazide;

30 2-Dibenzofurancarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

- 153 -

2-Dibenzofurancarboxylic acid, 2-[(5-fluoro-2-methylphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3,4-difluorophenyl)sulfonyl]-hydrazide;

5                   2-Dibenzofurancarboxylic acid, 2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-sulfonyl]hydrazide;

10                  2-Benzofurancarboxylic acid, 5-chloro-2-[(3-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3,4-difluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-chlorophenyl)-sulfonyl]hydrazide;

15                  2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-bromophenyl)-sulfonyl]hydrazide;

20                  2-Benzofurancarboxylic acid, 5-chloro-2-[(3-bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methylphenyl)-sulfonyl]hydrazide;

25                  2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chloro-2-methylphenyl)sulfonyl]hydrazide;

30                  2-Benzofurancarboxylic acid, 5-chloro-2-[(5-fluoro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

- 154 -

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

5           2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl)hydrazide;

10           2-Benzofurancarboxylic acid, 3,5-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

15           2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide;

20           2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(4-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(3-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-, 2-(phenylsulfonyl)hydrazide;

25           2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-, 2-(phenylsulfonyl)hydrazide;

30           Naphtho[2,3-b]furan-2-carboxylic acid, 3-methoxy-, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl)-sulfonyl hydrazide;

- 155 -

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-methylphenyl)-sulfonyl]hydrazide;

5                   2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-bromophenyl)-sulfonyl]hydrazide;

10                   2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethoxy)phenyl]sulfonyl]hydrazide;

15                   Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methoxy, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

20                   2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[2-(methylphenyl)-sulfonyl]hydrazide;

25                   2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chloro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[3-(bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chlorophenyl)-sulfonyl]hydrazide;

30                   2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)-hydrazide;

- 156 -

2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)-  
hydrazide;

5                   2-Benzofurancarboxylic acid, 5-chloro-3-methyl,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-fluorophenyl)sulfonyl]-  
hydrazide;

10                   2-Dibenzofurancarboxylic acid, 2-(2,6-difluorophenylsulfonyl)-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-bromophenyl)sulfonyl]-  
hydrazide;

15                   2-Dibenzofurancarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-methyl-3-nitrophenyl)-  
sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[2-chloro-5-(trifluoromethyl)-  
phenyl]sulfonyl]hydrazide;

20                   2-Dibenzofurancarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-bromo-2-  
trifluoromethoxyphenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]-hydrazide;

25                   2-Dibenzofurancarboxylic acid, 2-[(4-carboxyphenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4,5-dichloro-2-thienyl)-  
sulfonyl]hydrazide;

30                   2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-fluorophenyl)-  
sulfonyl]hydrazide;

- 157 -

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)sulfonyl]hydrazide;

10 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]-hydrazide;

15 2-Naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[[4-(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

20 2-Naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

25 2-Naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]-hydrazide;

30 3-Quinolinecarboxylic acid, 2-[(4-pentylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2(trifluoromethyl)phenyl]-sulfonyl]hydrazide;

- 158 -

3-Quinolinecarboxylic acid, 2-[[2,5-bis(2,2,2-trifluoroethoxyl)-phenyl]sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]-sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

- 159 -

Benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide;  
7-Ethoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-hydrazide;  
7-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)-hydrazide;  
2,3-Dihydro-benzofuran-5-carboxylic acid 2-(phenylsulfonyl)-

5 hydrazide;

5-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)-hydrazide;  
Isoquinoline-1-carboxylic acid 2-(phenylsulfonyl)hydrazide;  
1H-Indole-5-carboxylic acid 2-(phenylsulfonyl)hydrazide;

Quinoline-3-carboxylic acid 2-(phenylsulfonyl)hydrazide;

10 Quinoline-4-carboxylic acid 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-trifluoromethylphenyl)-

15 sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethylphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

20 1H-Indole-5-carboxylic acid, 2-[(3trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

25 6-Quinolinecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethylphenyl)sulfonyl]-  
hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

30 6-Quinolinecarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(4-fluoro-3-  
methylphenylsulfonyl)hydrazide;

- 160 -

7-Methoxy-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Methoxy-1H-indole-2-carboxylic acid, (phenylsulfonyl)hydrazide;

5-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5 5-Chloro-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
phenylsulfonyl]-hydrazide;

5-Methyl-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

and

10 5-Methyl-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
phenylsulfonyl]hydrazide.

7. A pharmaceutical composition comprising a compound of Claim 1 and a  
pharmaceutically acceptable carrier.

8. A method of treating or preventing neuronal loss associated with stroke,  
ischemia, CNS trauma, hypoglycemia or surgery, or treating a  
15 neurodegenerative disease, or treating or preventing the adverse consequences  
of the overstimulation of the excitatory amino acids, or treating anxiety,  
psychosis, glaucoma, CMV retinitis, diabetic retinopathy, urinary incontinence,  
migraine headache, convulsions, aminoglycoside antibiotics-induced hearing  
loss, Parkinson's disease, chronic pain, neuropathic pain, or inducing  
20 anesthesia, opioid tolerance or withdrawal, or enhancing cognition, comprising  
administering to a patient in need of such treatment an effective amount of a  
compound of Claim 1.



- 162 -

7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;  
2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)-

hydrazide;

5 Benzo[1,3] dioxole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1-Dibenzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-

10 hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-

hydrazide;

2-Dibenofurancarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-

15 hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(3,5-dichlorophenylsulfonyl)-

hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(3-chloro-2-methyl-

phenylsulfonyl)hydrazide;

20 Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-8-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(4-fluorophenyl)]-sulfonyl

hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-fluoro-

25 phenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-fluorophenyl)sulfonyl]-

hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-chloro-4-fluorophenyl)-

sulfonyl]hydrazide;

30 2-Dibenzofurancarboxylic acid, 2-[(3-cyanophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-cyanophenyl)-

sulfonyl]hydrazide;

- 163 -

2-Dibenzofurancarboxylic acid, 2-[(2,6-dichlorophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(2-chloro-

4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[[3,5-bis(trifluoromethyl)-phenyl]-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(5-fluoro-2-methylphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3,4-difluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3,4-difluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-bromophenyl)-sulfonyl]hydrazide;

- 164 -

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-methylphenyl)-sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

10 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chloro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(5-fluoro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

15 2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 6-methoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl)hydrazide;

25 2-Benzofurancarboxylic acid, 3,5-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

30 2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

- 165 -

2-Benzofurancarboxylic acid, 3,7-dimethoxy-,  
2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(4-fluorophenyl)-  
sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 3,7-dimethoxy-,  
2-[(3-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-,  
2-(phenylsulfonyl)hydrazide;

10 2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-,  
2-(phenylsulfonyl)hydrazide;

Naptho[2,3-b]furan-2-carboxylic acid, 3-methoxy-,  
2-(phenylsulfonyl)hydrazide;

15 2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl)-sulfonyl  
hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethyl)-  
phenyl]]sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-methylphenyl)-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro, 2-[(3-methylphenyl)  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro, 2-[(2-bromophenyl)-  
sulfonyl]hydrazide;

25 2-Benzofurancarboxylic acid, 5,7-dichloro, 2-[(3-bromophenyl)-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro, 2-[(3-chlorophenyl)-  
sulfonyl]hydrazide;

30 2-Benzofurancarboxylic acid, 5,7-dichloro,  
2-[[2-(trifluoromethoxy)phenyl]sulfonyl]hydrazide;

Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)-hydrazide;

- 166 -

2-Benzofurancarboxylic acid, 5-chloro-3-methoxy,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethyl)-  
phenyl]sulfonyl]hydrazide;

5        2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-methylphenyl)-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[2-(methylphenyl)-  
sulfonyl]hydrazide;

10       2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chloro-  
2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[3-(bromophenyl)-  
sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chlorophenyl)-  
sulfonyl]hydrazide;

15       2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethoxy)-  
phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)-hydrazide;

20       2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)-  
hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methyl,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)-hydrazide;

25       2-Dibenzofurancarboxylic acid, 2-[(2-fluorophenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-(2,6-difluorophenylsulfonyl)-  
hydrazide;

30       2-Dibenzofurancarboxylic acid, 2-[(2-bromophenyl)sulfonyl]-  
hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-  
hydrazide;

- 167 -

2-Dibenzofurancarboxylic acid, 2-[(4-methyl-3-nitrophenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[2-chloro-5-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

5 2-Dibenzofurancarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-bromo-2-trifluoromethoxyphenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]-hydrazide;

10 2-Dibenzofurancarboxylic acid, 2-[(4-carboxyphenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4,5-dichloro-2-thienyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)-sulfonyl]hydrazide;

15 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)-sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)sulfonyl]hydrazide;

25 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

30 2-Dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

- 168 -

2-Naphthalenecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[[2-(trifluoromethyl)phenyl]-  
sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[[4-(trifluoromethyl)phenyl]-  
sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)-  
sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-  
hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-(2-thienylsulfonyl)hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]-  
hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(4-pentylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2(trifluoromethyl)phenyl]-  
sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[[2,5-bis(2,2,2-trifluoroethoxyl)-  
phenyl]sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]-  
sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

- 169 -

6-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-hydrazide;

5 6-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide;

10 6-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

15 3-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

20 3-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

Benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

7-Ethoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)-

25 hydrazide;

5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

Isoquinoline-1-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-3-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

30 Quinoline-4-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

- 170 -

1H-Indole-5-carboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-trifluoromethylphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethylphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethylphenyl)sulfonyl]-

hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

6-Quinolinecarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(4-fluoro-3-  
methylphenylsulfonyl)hydrazide;

7-Methoxy-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Methoxy-1H-indole-2-carboxylic acid, hydrazide  
2-(phenylsulfonyl)hydrazide;

5-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Chloro-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
phenylsulfonyl]hydrazide;

5-Methyl-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;


5-Methyl-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-  
phenylsulfonyl]hydrazide; and

2-Naphthalenecarboxylic acid, 2-(phenylsulfonyl)hydrazide.

- 171 -

14. A method for inhibiting branched chain amino acid-dependent aminotransferases in a patient, the method comprising the step of administering to the patient a therapeutically effective amount of a compound of Claim 1.

15. The method of Claim 14, wherein Ar is selected from the group consisting of:



wherein  $R_6$  is is alkoxy, substituted alkoxy, halogen, aryloxy, substituted aryloxy, alkylamino, substituted alkylamino, arylalkylamino, substituted arylalkylamino, amide or substituted amide.

16. The method of Claim 14, wherein Ar is a bicyclic heteroaryl.

17. The method of Claim 14, wherein Ar is a tricyclic heteroaryl.

18. The method of Claim 14, wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ , and  $R_5$  are hydrogen.

19. The method of Claim 14, wherein the compound is:

Dibenzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

- 172 -

2,3,4,9-Tetrahydro-1H-carbazole-6-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

Benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5

7-Methoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Ethoxy-2-benzofurancarboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Chloro-benzofuran-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

2,3-Dihydro-benzofuran-5-carboxylic acid, 2-(phenylsulfonyl)-

10

hydrazide;

Benzo[1,3] dioxole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1-Dibenzofurancarboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

15

2-Dibenzofurancarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

2-Dibenofurancarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

20

2-Dibenzofurancarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(3,5-dichlorophenylsulfonyl)-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(3-chloro-2-methyl-phenylsulfonyl)hydrazide;

25

Quinoline-6-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-8-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(4-fluorophenyl)]-sulfonyl hydrazide;

30

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-fluoro-phenyl)sulfonyl]hydrazide;

- 173 -

2-Dibenzofurancarboxylic acid, 2-[(4-fluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-chloro-4-fluorophenyl)sulfonyl]hydrazide;

5 2-Dibenzofurancarboxylic acid, 2-[(3-cyanophenyl)sulfonyl]-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-cyanophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2,6-dichlorophenyl)sulfonyl]-hydrazide;

10 2-Benzofurancarboxylic acid, 5-bromo-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(2-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3,5-dichlorophenyl)sulfonyl]hydrazide;

15 2-Dibenzofurancarboxylic acid, 2-[[3,5-bis(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-nitro-, 2-(phenylsulfonyl)-hydrazide;

20 2-Dibenzofurancarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(5-fluoro-2-methylphenyl)sulfonyl]hydrazide;

25 2-Dibenzofurancarboxylic acid, 2-[(3,4-difluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

30 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-fluorophenyl)sulfonyl]hydrazide;

- 174 -

2-Benzofurancarboxylic acid, 5-chloro-2-[(3,4-difluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-chlorophenyl)-sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-bromophenyl)-sulfonyl]hydrazide;

10 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(2-methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methylphenyl)-sulfonyl]hydrazide;

15 2-Benzofurancarboxylic acid, 5-chloro-2-[(3-methoxyphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[(3-chloro-2-methylphenyl)sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 5-chloro-2-[(5-fluoro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

25 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

30 2-Benzofurancarboxylic acid, 6-methoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-methoxy-3-(1-methylethoxy), 2-(phenylsulfonyl)hydrazide;

- 175 -

2-Benzofurancarboxylic acid, 3,5-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

5 2-Benzofurancarboxylic acid, 4,6-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-(phenylsulfonyl)-hydrazide;

10 2-Benzofurancarboxylic acid, 3,7-dimethoxy-,  
2-[(2-trifluoromethylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-, 2-[(4-fluorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 3,7-dimethoxy-,  
2-[(3-methylphenyl)sulfonyl]hydrazide;

15 2-Benzofurancarboxylic acid, 3-methoxy-5-phenyl-,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 3-ethoxy-5-phenyl-,  
2-(phenylsulfonyl)hydrazide;

20 2-Benzofurancarboxylic acid, 3-methoxy-7-phenyl-,  
2-(phenylsulfonyl)hydrazide;

Naptho[2,3-b]furan-2-carboxylic acid, 3-methoxy-,  
2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-(phenylsulfonyl)-sulfonyl  
hydrazide;

25 2-Benzofurancarboxylic acid, 5,7-dichloro 2-[[2-(trifluoromethyl)-phenyl]]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-methylphenyl)-sulfonyl]hydrazide;

30 2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(2-bromophenyl)-sulfonyl]hydrazide;

- 176 -

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-bromophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5,7-dichloro 2-[(3-chlorophenyl)-sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 5,7-dichloro  
2-[[2-(trifluoromethoxy)phenyl]sulfonyl]hydrazide;

Benzofurancarboxylic acid, 5-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methoxy,  
2-(phenylsulfonyl)hydrazide;

10 2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethyl)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-methylphenyl)-sulfonyl]hydrazide;

15 2-Benzofurancarboxylic acid, 5-methyl-, 2-[2-(methylphenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chloro-2-methylphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[3-(bromophenyl)-sulfonyl]hydrazide;

20 2-Benzofurancarboxylic acid, 5-methyl-, 2-[(3-chlorophenyl)-sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 5-methyl-, 2-[[2-(trifluoromethoxy)-phenyl]sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methyl-, 2-(phenylsulfonyl)-hydrazide;

25 2-Benzofurancarboxylic acid, 4-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3-methyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 3,5-dimethyl-, 2-(phenylsulfonyl)-hydrazide;

2-Benzofurancarboxylic acid, 5-chloro-3-methyl,

30 2-(phenylsulfonyl)hydrazide;

2-Benzofurancarboxylic acid, 5-amino-, 2-(phenylsulfonyl)-hydrazide;

- 177 -

2-Dibenzofurancarboxylic acid, 2-[(2-fluorophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-(2,6-difluorophenylsulfonyl)-hydrazide;

5                   2-Dibenzofurancarboxylic acid, 2-[(2-bromophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

10                   2-Dibenzofurancarboxylic acid, 2-[(4-methyl-3-nitrophenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[2-chloro-5-(trifluoromethyl)phenyl]sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-trifluoromethoxyphenyl)sulfonyl]hydrazide;

15                   2-Dibenzofurancarboxylic acid, 2-[(4-bromo-2-trifluoromethoxyphenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2-nitrophenyl)sulfonyl]-hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(4-carboxyphenyl)sulfonyl]-hydrazide;

20                   2-Dibenzofurancarboxylic acid, 2-[(4,5-dichloro-2-thienyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chlorophenyl)sulfonyl]hydrazide;

25                   2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-fluorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

30                   2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-chloro-2-methoxyphenyl)sulfonyl]hydrazide;

- 178 -

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(5-bromo-2-methoxyphenyl)sulfonyl]hydrazide;

2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(4-ethylphenyl)sulfonyl]hydrazide;

5 2-Benzofurancarboxylic acid, 7-methoxy-, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

2-Dibenzofurancarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

10 2-Naphthalenecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2-(trifluoromethyl)phenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-(trifluoromethyl)phenyl)sulfonyl]hydrazide;

15 2-Naphthalenecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

20 2-Naphthalenecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

2-Naphthalenecarboxylic acid, 2-[(2,3-dichlorophenyl)sulfonyl]hydrazide;

2-Naphthalenecarboxylic acid, 2-[(4-ethylphenyl)sulfonyl]-hydrazide;

25 3-Quinolinecarboxylic acid, 2-[(4-pentylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2(trifluoromethyl)phenyl]sulfonyl]hydrazide;

3-Quinolinecarboxylic acid, 2-[[2,5-bis(2,2,2-trifluoroethoxy)phenyl]sulfonyl]hydrazide;

30 3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

3-Quinolinecarboxylic acid, 2-[[2-(trifluoromethoxy)phenyl]sulfonyl]hydrazide;

- 179 -

3-Quinolinecarboxylic acid, 2-[(4-methylphenyl)sulfonyl]-hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;  
3-Quinolinecarboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;  
5 3-Quinolinecarboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;  
3-Quinolinecarboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;  
6-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-  
hydrazide;  
10 3-Quinolinecarboxylic acid, 2-[(2-trifluoromethylphenyl)sulfonyl]-  
hydrazide;  
6-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;  
3-Quinolinecarboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
15 sulfonyl]hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-  
hydrazide;  
3-Quinolinecarboxylic acid, 2-[(3,5-dichlorophenyl)sulfonyl]-  
hydrazide;  
20 6-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-  
hydrazide;  
3-Quinolinecarboxylic acid, 2-[(3-chloro-4-fluorophenyl)sulfonyl]-  
hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;  
25 3-Quinolinecarboxylic acid, 2-[(3-methoxyphenyl)sulfonyl]-hydrazide;  
6-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;  
3-Quinolinecarboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;  
Benzofuran-2-carboxylic acid 2-(phenylsulfonyl)hydrazide;  
7-Ethoxy-2-benzofurancarboxylic acid 2-(phenylsulfonyl)-hydrazide;  
30 7-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)-hydrazide;  
2,3-Dihydro-benzofuran-5-carboxylic acid 2-(phenylsulfonyl)-  
hydrazide;

- 180 -

5-Chloro-benzofuran-2-carboxylic acid 2-(phenylsulfonyl)-hydrazide;

Isoquinoline-1-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

Quinoline-3-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5 Quinoline-4-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-chlorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-fluorophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-trifluoromethylphenyl)-

10 sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethylphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

15 1H-Indole-5-carboxylic acid, 2-[(3trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-bromophenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(3-methylphenyl)sulfonyl]-hydrazide;

1H-Indole-5-carboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

20 6-Quinolinecarboxylic acid, 2-[(2-chlorophenyl)sulfonyl]-hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethylphenyl)sulfonyl]-  
hydrazide;

6-Quinolinecarboxylic acid, 2-[(3-trifluoromethoxyphenyl)-  
sulfonyl]hydrazide;

25 6-Quinolinecarboxylic acid, 2-[(2-methylphenyl)sulfonyl]-hydrazide;

Dibenzofuran-2-carboxylic acid, 2-(4-fluoro-3-  
methylphenylsulfonyl)hydrazide;

7-Methoxy-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

7-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

30 5-Methoxy-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)hydrazide;

5-Chloro-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

- 181 -


5-Chloro-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-phenylsulfonyl]hydrazide;

5-Methyl-1H-indole-2-carboxylic acid, 2-(phenylsulfonyl)-hydrazide;

5-Methyl-1H-indole-2-carboxylic acid, 2-[(2-trifluoromethyl)-phenylsulfonyl]hydrazide; and

2-Naphthalenecarboxylic acid, 2-(phenylsulfonyl)hydrazide.


20. A method for preparing a compound of Formula I



wherein:

R<sub>3</sub> is H, or F, Br, alkyl, carboxy, alkoxy, substituted alkoxy,

R<sub>1</sub>, R<sub>2</sub>, R<sub>4</sub>, and R<sub>5</sub> are independently, H, halogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cyano, nitro, amino, alkylamine, and thioalkyl, Ar is bicyclic heteroaryl, tricyclic heteroaryl, substituted bicyclic heteroaryl, except (6-methoxybenzofuran)-2-nyl, 3-quinolinyl which comprises reacting a hydrazide of the formula ArC(O)NH<sub>2</sub>NH<sub>2</sub> with a sulfonyl halide of the formula:



where R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and Ar are as defined above and Hal is halogen.

- 182 -

21. Use of a compound of Formula I as claimed in claim 1 for preparation of a medicament for treatment or or preventing neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycemia or surgery, or treating a neurodegenerative disease, or treating or preventing the adverse consequences of the overstimulation of the excitatory amino acids, or treating anxiety, psychosis, glaucoma, CMV retinitis, diabetic retinopathy, urinary incontinence, migraine headache, convulsions, aminoglycoside antibiotics-induced hearing loss, Parkinson's disease, chronic pain, neuropathic pain, or inducing anesthesia, opioid tolerance or withdrawal, or enhancing cognition.

22. A medicine for treatment or or preventing neuronal loss associated with stroke, ischemia, CNS trauma, hypoglycemia or surgery, or treating a neurodegenerative disease, or treating or preventing the adverse consequences of the overstimulation of the excitatory amino acids, or treating anxiety, psychosis, glaucoma, CMV retinitis, diabetic retinopathy, urinary incontinence, migraine headache, convulsions, aminoglycoside antibiotics-induced hearing loss, Parkinson's disease, chronic pain, neuropathic pain, or inducing anesthesia, opioid tolerance or withdrawal, or enhancing cognition comprising a therapeutically effective amount of a compound as claimed in claim 1.